Appendix 4

Effects of Temperature, Dissolved Oxygen/Total Dissolved Gas, Ammonia, and pH on Salmonids

Implications for California's North Coast TMDLs

Katharine Carter
Environmental Scientist
North Coast Regional Water Quality Control Board

TABLE OF CONTENTS

CHAPTER	1: TEMPERATURE	1
1.1 Introdu	ction	1
1.2 Temper	ature Metrics	1
1.3 Adult M	ligration and Holding	3
1.3.1	Steelhead Trout Migration	
1.3.2	Chinook Salmon Migration and Holding	4
1.3.3	Coho Salmon Migration	
1.4 Spawnii	ng, Incubation, and Emergence	7
1.4.1	Steelhead Spawning, Incubation, and Emergence	7
1.4.2	Chinook Spawning, Incubation, and Emergence	9
1.4.3	Coho Spawning, Incubation, and Emergence	10
1.5 Freshwa	ater Rearing and Growth	
1.5.1	Steelhead Freshwater Rearing and Growth	12
1.5.2	Chinook Freshwater Rearing and Growth	14
1.5.3	Coho Freshwater Rearing and Growth	16
1.6 Lethalit	y	18
1.6.1	Steelhead Lethality	18
1.6.2	Chinook Lethality	18
1.6.3	Coho Lethality	18
1.7 Disease.		19
1.7.1	Ichthyophthiriasis (Ich)	22
1.7.2	Ceratomyxosis	22
1.7.3	Columnaris	23
1.8 TMDL 7	Femperature Thresholds	24
	-	
CHAPTER 2	2: DISSOLVED OXYGEN AND TOTAL DISSOLVED GAS	26
2.1 Introduc	ction	26
2.2 Effects of	of Low Dissolved Oxygen Concentrations on Salmonids	26
2.2.1	Adult Migration	
2.2.2	Incubation/Emergence	26
2.2.3	Incubation mortality	27
2.2.4	Incubation growth	28
2.2.5	Incubation avoidance/preference	29
2.2.6	Emergence mortality	29
2.2.7	Freshwater Rearing and Growth	
	2.2.7.1 Swimming and activity	30
	2.2.7.2 Growth	30
	2.2.7.3 Avoidance and preference	32
2.2.8	Lethality	
2.3 Effects of	of High Total Dissolved Gas Concentrations on Salmonids	
	-	
CHAPTER 3	3: AMMONIA	35
	ction	

3.2 Ammonia Speciation	35
3.3 Ammonia Toxicity	
3.4 Ammonia Criteria	
CHAPTER 4: pH	39
4.1 Introduction	39
4.2 Effects of High pH	39
4.3 Effects of Low pH	
•	
REFERENCES	42

LIST OF TABLES

Table 1: Effects of Temperature in Considering Adult Steelhead and Migration	4
Table 2: Effects of Temperature in Considering Adult Chinook and Migration and	
Holding	5
Table 3: Effects of Temperature in Considering Adult Coho and Migration	
Table 4: Effects of Temperature in Considering Steelhead Incubation and Emergence	
Table 5: Effects of Temperature in Considering Steelhead, Chinook, and Coho	
Spawning	8
Table 6: Effects of Temperature in Considering Chinook Incubation and Emergence	9
Table 7: Effects of Temperature in Considering Coho Incubation and Emergence	
Table 8: Effects of Temperature in Considering Juvenile Steelhead Rearing and	
Growth	13
Table 9: Effects of Temperature in Considering Juvenile Chinook Rearing and	
Growth	15
Table 10: Effects of Temperature in Considering Juvenile Coho Rearing and	
Growth	17
Table 11: Effects of Temperature in Considering Lethality and Salmonids	19
Table 12: Effects of Temperature in Considering Disease and Salmonids	20-21
Table 13: Life Stage Temperature Thresholds	
Table 14: Lethal Temperature Thresholds	25
Table 15: Dissolved oxygen concentrations and their effects salmonid embryo and	
larval stages	29
Table 16: Dissolved oxygen concentrations and their effects on salmonid life stages other	
than embryonic and larval	33
Table 17: pH-Dependent Values of the Criterion Maximum Concentration (CMC) of	
Total Ammonia as Nitrogen (mg N/L) in Freshwater when Salmonids are	
Present	37
Table 18: Temperature and pH-Dependent Values of the Criterion Continuous	
Continuation (CCC) for Total Ammonia as Nitrogen (mg N/L) in Freshwater	
when Fish Early Life Stages are Present	37
Table 19: Temperature and pH-Dependent Values of the Criterion Continuous	
Continuation (CCC) for Total Ammonia as Nitrogen (mg N/L) in Freshwater	
when Fish Early Life Stages are Absent	38
Table 20: Reactions of 10 rainbow trout to various pH levels during gradual acclimation	
experiments (0 .2 to 0.4 of a pH unit/day)	40
LIST OF FIGURES	
Figure 1: Chemical Speciation of Ammonia	35

CHAPTER 1. TEMPERATURE

1.1 Introduction

Temperature is one of the most important environmental influences on salmonid biology. Most aquatic organisms, including salmon and steelhead, are poikilotherms, meaning their temperature and metabolism is determined by the ambient temperature of water. Temperature therefore influences growth and feeding rates, metabolism, development of embryos and alevins, timing of life history events such as upstream migration, spawning, freshwater rearing, and seaward migration, and the availability of food. Temperature changes can also cause stress and lethality (Ligon et al. 1999). Temperatures at sublethal levels can effectively block migration, lead to reduced growth, stress fish, affect reproduction, inhibit smoltification, create disease problems, and alter competitive dominance (Elliott 1981, USEPA 1999a). Further, the stressful impacts of water temperatures on salmonids are cumulative and positively correlated to the duration and severity of exposure. The longer the salmonid is exposed to thermal stress, the less chance it has for long-term survival (Ligon et al. 1999).

A literature review was performed to evaluate temperature needs for the various life stages of steelhead trout (*Oncorhynchus mykiss*), coho salmon (*Oncorhynchus kisutch*), and Chinook salmon (*Oncorhynchus tschawytscha*). The purpose of this review was to identify temperature thresholds that are protective of salmonids by life stage, as a basis for evaluating stream temperatures in California temperature TMDLs within the North Coast region.

This review included USEPA temperature guidance, Oregons' and Washingtons' temperature standards reviews, reports that compiled and summarized existing scientific information, and laboratory and field studies. When possible, species-specific needs were summarized by the following life stages: migrating adults, spawning and incubation/emergence, and freshwater rearing and growth. Additionally, the effects of temperature on disease and lethality are also discussed. Some of the references reviewed covered salmonids as a general class of fish, while others were species specific. Information for fall run coho salmon, spring/summer, fall, and winter steelhead, and spring and fall run Chinook salmon are compiled by life stage in Table 1 through Table 12.

1.2 Temperature Metrics

In considering the effect of temperature on salmonids, it is useful to have a measure of chronic and acute (i.e. sub-lethal and lethal) temperature exposures. A common measure of chronic exposure is the maximum weekly average temperature (MWAT). The MWAT is the maximum seasonal or yearly value of the mathematical mean of multiple, equally spaced, daily temperatures over a running seven-day consecutive period (Brungs and Jones 1977, p.10). In other words, it is the highest single value of the seven-day moving average temperature. A common measure of acute effects is the instantaneous maximum. A third metric, the maximum weekly maximum temperature (MWMT), can be used as a

measure of both chronic and acute effects. The MWMT is also known as the seven-day average of the daily maximum temperatures (7-DADM), and is the maximum seasonal or yearly-value of the daily maximum temperatures over a running seven-day consecutive period. The MWMT is useful because it describes the maximum temperatures in a stream, but is not overly influenced by the maximum temperature of a single day.

Much of the information reported in the literature characterizes temperature needs with terms such as "preferred" or "optimum". Preferred stream temperatures are those that fish most frequently inhabit when allowed to freely select temperatures in a thermal gradient (USEPA 1999a). An optimum range provides suitable temperatures for feeding activity, normal physiological response, and normal behavior (without symptoms of thermal stress) (USEPA 1999a). Optimal temperatures have also been described as those temperatures at which growth rates, expressed as weight gain per unit of time, are maximal for the life stage (Armour 1991).

Salmonid stocks do not tend to vary much in their life history thermal needs, regardless of their geographic location. In the 2001 USEPA document, *Summary of Technical Literature Examining the Physiological Effects of Temperature on Salmonids*, the case is made that there is not enough significant genetic variation among stocks or among species of salmonids to warrant geographically specific water temperature standards.

Climate conditions vary substantially among regions of the State and the entire Pacific Northwest. ...Such [varying climatic] conditions could potentially have led to evolutionary adaptations, resulting in development of subspecies differences in thermal tolerance. ...[However,] the literature on genetic variation in thermal effects indicates occasionally significant but very small differences among stocks and increasing differences among subspecies, species, and families of fishes. Many differences that had been attributed in the literature to stock differences are now considered to be statistical problems in analysis, fish behavioral responses under test conditions, or allowing insufficient time for fish to shift from field conditions to test conditions (Mathur & Silver 1980, Konecki et al. 1993, both as cited in USEPA 2001).

Additionally:

There are many possible explanations why salmonids have not made a significant adaptation to high temperature in streams of the Pacific Northwest. Temperature tolerance is probably controlled by multiple genes, and consequently would be a core characteristic of the species not easily modified through evolutionary change without a radical shift in associated physiological systems. Also, the majority of the life cycle of salmon and steelhead is spent in the ocean rearing phase, where the smolt, subadults, and adults seek waters with temperatures less than 59°F (15°C) (Welch et al, 1995, as cited in USEPA 2001).

As a result, literature on the temperature needs of coho and Chinook salmon and steelhead trout stemming from data collected in streams outside Northern California are cited in this document and are considered relevant to characterizing the thermal needs of salmonids, which use Northern California rivers and streams.

1.3 Adult Migration and Holding

All of the adult migration and holding temperature needs referenced in this section can be found in Table 1 through Table 3. Salmon and trout respond to temperatures during their upstream migration (Bjornn and Reiser 1991). Delays in migration have been observed in response to temperatures that were either too cold or too warm. Most salmonids have evolved with the temperature regime they historically used for migration and spawning, and deviations from the normal pattern can affect survival (Spence et al. 1996).

In a 2003 USEPA document entitled *EPA Region 10 Guidance for Pacific Northwest State and Tribal Water Quality Standards*, it is recommended that the 7-DADM should not exceed 18°C in waters where both adult salmonid migration and "non-core" juvenile rearing occur during the period of summer maximum temperatures. The document does not define what constitutes the "summer" period. Non-core juvenile rearing is defined as moderate to low density salmon and trout rearing usually occurring in the mid or lower part of the basin, as opposed to areas of high density rearing which are termed "core" rearing areas. This criterion is derived from analysis and synthesis of past laboratory and field research. The USEPA believes that this temperature recommendation will protect against lethal conditions, prevent migration blockage, provide optimal or near optimal juvenile growth conditions, and prevent high disease risk by minimizing the exposure time to temperatures which can lead to elevated disease rates.

A 7-DADM temperature of 20°C is recommended by the USEPA (2003) for waterbodies that are used almost exclusively for migration during the period of summer maximum temperatures.

EPA believes that a 20°C criterion would protect migrating juveniles and adults from lethal temperatures and would prevent migration blockage conditions. However, EPA is concerned that rivers with significant hydrologic alterations (e.g., rivers with dams and reservoirs, water withdrawals, and /or significant river channelization) may experience a loss of temperature diversity in the river, such that maximum temperatures occur for an extended period of time and there is little cold water refugia available for fish to escape maximum temperatures. In this case, even if the river meets a 20°C criterion for maximum temperatures, the duration of exposure to 20°C temperatures may cause adverse effects in the form of increased disease and decreased swimming performance in adults, and increased disease, impaired smoltification, reduced growth, and increased predation for late emigrating juveniles...(USEPA 2003).

Therefore, the USEPA recommends a narrative provision to protect and, if possible, restore the natural thermal regime accompany the 7-DADM 20°C criterion for rivers with significant hydrologic alterations.

In an exhaustive study of both laboratory and field studies of temperature effects on salmonids and related species, USEPA (1999a, 2001) concluded that temperatures of approximately 22-24°C limit salmonid distribution, i.e., they totally eliminate salmonids from a location. USEPA (1999a) also notes that changes in competitive interactions between fish species can lead to a transition in dominance from salmonids to other species at temperatures 2-4°C lower than the range of total elimination.

1.3.1 Steelhead Trout Migration

In a 2002 review of numerous studies, Washington State Department of Ecology (WDOE) concluded that daily average temperatures of 21-24°C are associated with avoidance behavior and migration blockage in steelhead trout. WDOE suggests that the MWMT should not exceed 17-18°C, and daily maximum temperatures should not exceed 21-22°C to be fully protective of adult steelhead migration.

Table 1: Effects of Temperature in Considering Adult Steelhead and Migration

С	_		
24 23 22	21-24 Average daily temperature associated with avoidance and	22-24 Temperature range which eliminates salmonids	s from an area (3,4)
21	migration blockage (2)	21-22 Daily maximum temperature should not exceed this to be fully protective (2)	18-22 Temperature
20	20 MWMT should not exceed this Should be used in conjunction wit natural thermal regime for r	range at which transition in dominance from	
19			salmonids to other
18	17-18 MWMT should not exceed this to be fully protective (2)	18 MWMT should not exceed this where migration and non-core rearing occur (1)	species occurs (4)
17	uns to be runy protective (2)		

Sources: (1) USEPA 2003, (2) WDOE 2002, (3) USEPA 2001, (4) USEPA 1999a

1.3.2 Chinook Salmon Migration and Holding

USEPA (2001) cited various literature sources that identified thermal blockages to Chinook salmon migration at temperatures ranging from 19-23.9°C, with the majority of references citing migration barriers at temperatures around 21°C.

A radio tracking study on spring Chinook revealed that when maximum temperatures of 21.1°C were reached, a thermal barrier to migration was established (Bumgarner et al. 1997, as cited by USEPA 1999a). Bell (1986) reviewed various studies and notes spring Chinook migrate at water temperatures ranging from 3.3-13.3°C, while fall Chinook migrate at temperatures of 10.6-19.6°C. Preferred temperatures for Chinook range from 7.2-14.5°C (Bell 1986). Based on a technical literature review, WDOE (2002) concluded that daily maximum temperatures should not exceed 21-22°C during Chinook migration.

Table 2: Effects of Temperature in Considering Adult Chinook and Migration and Holding

	ie 2. Effects of Temperature in Consider	ering Adult Chinook and Migrati	on and Holding	1	
°C		MIGRATION			
23	23 Klamath Basin fall Chinook begin migration upstream at temperatures as high as 23C if temperatures are rapidly falling (6) 22 Klamath Basin fall Chinook will not migrate upstream when mean daily temperatures are 22C or greater (6)	22-24 Temperature range which eliminates salmonids from an area (3,5)			
21	21-22 Daily maximum temperature should not exceed this range to be protective of migration (2)	21 Most references cite as thermal block to migration (3) 21 Klamath Basin fall Chinook will not migrate upstream if temperatures are 21C or above and rising (6)	19-23.9 Range of temperatures causing thermal blockage to migration (3)	18-22 Temperature range at which transition in dominance	
20	20 MWMT should not exceed this in wa for migration. Should be used in conju about protecting/restoring the natural significant hydrologic	nction with a narrative provision thermal regime for rivers with		from salmonids to other species occurs (5)	
19					
18			18 MWMT should where migration rearing of	and non-core	
17	16-17 MWMT should be below this		J	· /	
16	where Chinook are holding (2)	10.6-19.6 Temperature range			
15		where adult fall Chinook migrate	10.11	.,	
14		(4)	13-14 Average daily temperature should be below this where spring Chinook are holding (2)		
13					
12 11	7.2-14.5 Preferred temperatures for Chinook (4)				
10	C.I.I.SOR (1)		3.3-13.3 Temperat		
9 8 7 6			addit spring clim	our migrate (1)	
5					
4					
3					

Sources: (1) USEPA 2003, (2) WDOE 2002, (3) USEPA 2001, (4) Bell 1986, (5) USEPA 1999a, (6) Strange 2006

Utilizing radio telemetry to track the movements and monitor the internal body temperatures of adult fall Chinook salmon during their upriver spawning migration in the Klamath basin, Strange (2006) found that fall Chinook will not migrate upstream when mean daily temperatures are $\geq 22^{\circ}$ C. Strange also noted that adult fall Chinook in the

Klamath basin will not migrate upstream if temperatures are 21°C or above and rising, but will migrate at temperatures as high as 23°C if temperatures are rapidly falling.

Spring Chinook begin entering freshwater streams during a relatively cool-water season but must hold throughout the warm summer period, awaiting cooler spawning temperatures (ODEQ 1995a). The cumulative effects of management practices such as elevated water temperatures, reduced cover from large woody debris, and reduced resting pool area due to pool filling increase the susceptibility of holding adult fish to mortality from thermal effects (The Oregon Department of Environmental Quality [ODEQ] 1995a). WDOE states that where spring Chinook are holding over for the summer prior to spawning the average daily water temperature should be below 13-14°C and the MWMT should be below 16-17°C (WDOE 2002).

1.3.3 Coho Salmon Migration

Migration for coho is delayed when water temperatures reach 21.1°C, and the preferred water temperatures for coho range from 11.7-14.5°C (Bell 1986). In California coho salmon typically migrate upstream when water temperatures range from 4-14°C (Briggs, 1953 and Shapovalov and Taft, 1954, as cited by Hassler, 1987). WDOE reviewed various studies and concluded that to be protective of adult coho migration, MWMTs should not exceed 16.5°C (WDOE 2002).

Table 3: Effects of Temperature in Considering Adult Coho and Migration

	le 3. Effects of Temperature in Con		18.44.01.
°C		MIGRATION	
24 23 22	22-24 Temperature range which el area (3,6		
21	21.1 Migration is delayed when tem	peratures reach this value (4)	18-22 Temperature range at which
20	20 MWMT should not exceed this exclusively for migration. Should be narrative provision about protecting regime for rivers with significant	transition in dominance from salmonids to other species occurs (6)	
19			
18	18 MWMT should not exceed this v rearing occu		
17			
16	16.5 MWMT sho	ould not exceed this value to be	fully protective (2)
15			
14			
13	11.7-14.5 Preferred temperature		
12	range (4)	4-14 Temperature range at v	which migration typically occurs (5)
11	11.4 Preferred temperature (7)		

Sources: (1) USEPA 2003, (2) WDOE 2002, (3) USEPA 2001, (4) Bell 1986, (5) Briggs 1953, Shapovalov and Taft 1954, as cited by Hassler 1987, (6) USEPA 1999a, (7) Reutter and Herdendorf 1974

1.4 Spawning, Incubation, and Emergence

All of the spawning, incubation, and emergence temperature needs referenced in this section can be found in Table 4 through Table 7. Many sources have stated that temperature affects the time of migration i adults and thus the time of spawning, which influences the incubation temperature regime, which in turn influences survival rates, development rates, and growth of embryos and alevins (Murray and McPhail 1988). USEPA Region 10 recommends that the 7-DADM temperatures should not exceed 13°C for salmonid spawning, egg incubation, and fry emergence (USEPA 2003). Optimum temperatures for salmonid egg survival ranges from 6-10°C (USEPA 2001).

1.4.1 Steelhead Spawning, Incubation, and Emergence

In a discussion paper and literature summary evaluating temperature criteria for fish species including salmonids and trout, WDOE (2002) cites studies showing that steelhead were observed spawning in temperatures ranging from 3.9-21.1 °C, and that the preferred temperatures for steelhead spawning range from 4.4-12.8 °C. In a review of various studies, Bell (1986) concludes that steelhead spawning occurs at water temperatures ranging from 3.9-9.4 °C.

Steelhead and rainbow trout eggs had the highest survival rates between 5-10°C according to Myrick and Cech (2001), and while they can tolerate temperatures as low as 2°C or as high as 15°C, mortality is increased at these temperatures. WDOE (2002) reviewed literature on the survival of steelhead and rainbow trout embryos and alevins at various temperatures and concluded that the average water temperature should not exceed 7-10°C throughout development, and the maximum daily average temperature should be below 11-12°C at the time of hatching (WDOE 2002).

Table 4: Effects of Temperature in Considering Steelhead Incubation and Emergence

°C	INCUBATION AND EMERGENCE					
15	15 Steelhead and rainbow trout eggs can survive at temperatures as high as this but mortality is high compared to lower temperatures (3)					
14						
13	13 MWMT should not exce	eed this value to be protective emergence (1)	of spawning, egg incubation, and fry			
12 11	11-12 Maximum daily average	ge temperature should be below	w this range at the time of hatching (2)			
10 9 8 7 6	5-10 Steelhead and rainbow trout eggs had the highest survival within this range (3)	6-10 Optimum temperature for salmonid eggs survival to hatching (4)	7-10 Average daily temperature should not exceed this range throughout embryo development (2)			
5						
4						
2	2 Steelhead and rainbow trout eggs can survive at temperatures as low as this but mortality is high compared to higher temperatures (3)					

Sources: (1) USEPA 2003, (2) WDOE 2002, (3) Myrick and Cech 2001, (4) USEPA 2001

Table 5: Effects of Temperature in Considering Steelhead, Chinook, and Coho Spawning

°C	ie 3. Elicets o	Steelhead	n considering s		Chinook		Col	ho	All Salmonids
21 20									
19 18									
17 16									
15									
14	3.9-21.2 Steelhead			13-15.5 Temp. range at which pre-spawning mortality becomes pronounced in	of refs. cite daily max temps. associated with spawning below this level (2)	5.6-17.7 Range of			
13	observed spawning in this temp. range (2)			ripe spring Chinook (4)		temps. associated with spawning from	13 Daily maximum temp. not to exceed this value to be protective (6)		13 MWMT not exceed this value during spawning, egg incubation, and fry emergence (1)
12 11					5.6-13.9	references			
11					Recommended	reviewed (2)	10 MWAT not	4.4-13.3	
10			4.4-12.8 Preferred	5.6-12.8 Recommended	temperature range for spawning (5)		exceed this value to be protective (6)	Typical temps. during which spawning occurs (2)	
9			temp. range for spawning	temperature range for			4.5-9.4		
8 7		3.9-9.4 Temp.	(2)	spawning (4)			Preferred		
6		range where					spawning		
5		spawning occurs (3)					temperature range (3)		
4		occurs (3)					range (3)		
3									

Sources: (1) USEPA 2003, (2) WDOE 2002, (3) Bell 1986, (4) ODEQ 1995a, (5) Reiser and Bjornn 1979 as cited by Armour et al. 1991, (6) Brungs and Jones 1977

1.4.2 Chinook Spawning, Incubation, and Emergence

ODEQ (1995a) reviewed numerous studies and recommended a temperature range of 5.6-12.8°C for spawning Chinook. A discussion paper and literature summary by WDOE in 2002 found that the literature reviewed noted a wide range of temperatures associated with Chinook spawning (5.6-17.7°C), although the majority of these temperature observations cite daily maximum temperatures below 14.5°C. A spawning temperature range of 5.6-13.9°C is recommended for spring, summer, and fall Chinook salmon populations in the Pacific Northwest (Reiser and Bjornn 1979, as cited by Armour et al. 1991). When ripe adult spring Chinook females experience temperatures above 13-15.5°C, pre-spawning adult mortality becomes pronounced (ODEQ 1995a). Additionally, there is decreased survival of eggs to the eyed stage and alevin development is inhibited due to the exposure of the ripe female to warm temperatures, even if the stream temperatures during the egg and alevin development are appropriate (ODEQ 1995a).

Table 6: Effects of Temperature in Considering Chinook Incubation and Emergence

°C	INCUBATION AND EMERGENCE						
20 19 18 17	17.5-20 The hi	ghest single day maximum temp		exceed this range to p	rotect eggs and	l embryos	
16							
15			1		ı		
14		13.5-14.5 Daily maximum	14 Moderate e	mbryo survival (6)	1		
13		temperatures should not exceed this from fertilization through initial fry development (5)	13 MWMT should not exceed this value to be protective of spawning, egg incubation, and fry emergence (1)				
12				11-12.8 Average		17167	
11	5-14.4 Recommended temp.	11 High embryo survival (6)		daily temperatures should be below this range at beginning of incubation (2)	2-14 Range of temps.	1.7-16.7 Eggs can survive these temps.	
9	range for incubation (4)	9-10 Optimal temp. should be below this range (5) 8-9 Seasonal ave. temps.	4-12 Lowest levels of egg mortality at	6-10 Optimum	for normal embryo develop-	but mortality is greatly increased	
8		should not exceed this range from fertilization through initial fry development (2) 8 High embryo survival (6)	these temps. (3)	temperature for salmonid eggs survival to hatching (5)	ment (6)	at the extremes (3)	
7							
6							
5		5 High embryo survival (6)					
4							
3							
2		2 Poor embryo su	ırvival (6)				
1							

Sources: (1) USEPA 2003, (2) WDOE 2002, (3) Myrick and Cech 2001, (4) Reiser and Bjornn 1979, as cited by Armour et al. 1991, (5) USEPA 2001, (6) Murray and McPhail 1988

WDOE (2002) reviewed numerous references on the effects of various temperatures on Chinook incubation and development and used these studies to derive the temperatures that are protective of Chinook salmon from fertilization through fry development. These reviewed references include laboratory studies assessing Chinook embryo survival at various constant temperatures, studies attempting to mimic naturally fluctuating temperatures experienced by incubating eggs, studies which have made stepwise reductions in the incubation temperatures as incubation progressed to evaluate survival of eggs, and studies on the effects of transferring eggs to optimal constant incubation temperatures after they had been exposed to higher temperatures for various periods. As a result of this review, WDOE (2002) recommends that average daily temperatures remain below 11-12.8°C at the initiation of incubation, and that the seasonal average should not exceed 8-9°C in order to provide full protection from fertilization through initial fry development. The highest single day maximum temperature should not exceed 17.5-20°C to protect eggs and embryos from acute lethal conditions (WDOE 2002).

USEPA (2001) reviewed multiple literature sources and concluded that optimal protection from fertilization through initial fry development requires that temperatures be maintained below 9-10°C, and that daily maximum temperatures should not exceed 13.5-14.5°C. Reiser and Bjornn (1979, as cited by Armour et al. 1991) recommended temperatures of 5.0-14.4°C for spring, summer and fall Chinook salmon incubation in the Pacific Northwest. Myrick and Cech (2001) reviewed studies on the Sacramento-San Joaquin River and concluded that the lowest levels of Chinook egg mortality occurred at temperatures between 4-12°C, and while eggs can survive at temperatures from 1.7-16.7°C, mortality is greatly increased at the temperature extremes.

Embryo survival was studied in a laboratory experiment conducted by Murray and McPhail (1988). They incubated five species of Pacific salmon, including Chinook, at five incubation temperatures (2, 5, 8, 11, 14°C). Chinook embryo survival was high at 5, 8, and 11°C, but survival was moderate at 14°C and poor at 2°C. As a result of their study, Murray and McPhail concluded that the range of temperatures for normal embryo development is > 2°C and <14°C (Murray and McPhail 1988).

1.4.3 Coho Spawning, Incubation, and Emergence

In 2002, WDOE found that several studies and literature reviews state that spawning activity in coho may typically occur in the range of 4.4-13.3°C. According to a review by Bell (1986), preferred spawning temperatures range from 4.5-9.4°C. Brungs and Jones (1977) used existing data on the optimum and range of temperatures for coho spawning and embryo survival to create criteria using protocols from the National Academy of Sciences and National Academy of Engineering. The resultant criteria were that the MWAT should not exceed 10°C and the daily maximum temperature should not exceed 13°C to be protective of coho (Brungs and Jones 1977, p.16).

In a discussion paper and literature summary WDOE (2002) reviewed studies that assessed the survival of embryos and alevin at various temperatures. Based on the findings of these studies WDOE (2002) has determined that the average daily temperature during the incubation period should be at or below 8-10°C to fully support this coho salmon life stage. According to a review of various literature sources by Bell

(1986), the preferred emergence temperatures for coho range from 4.5-13.3°C. USEPA (2001) concluded that to fully support pre-emergent stages of coho development MWMTs should not exceed 9-12°C.

Table 7: Effects of Temperature in Considering Coho Incubation and Emergence

°C	INCUBATION AND EMERGENCE					
14		14 Upper	limit for normal e	mbryo d	evelopment (5)	
	13 MWMT sl	hould not exceed thi	s value to be		aily maximum temperature	
13	protective of s	pawning, egg incub	ation, and fry	should	d not exceed this value to be	
		emergence (1)			protective (6)	
12						
11			9-12 MWMT s	hould		
10	6-10 Optimum temperature for	8-10 Ave. daily temp. during incubation	not exceed this r be fully protect	ange to	10 MWAT should not exceed this to be protective (6)	4.5-13.3 Preferred emergence
9	salmonid eggs	should be at or				temperature
8	survival to	below this to be supportive (2)				range (3)
7	hatching (4)					
6						
5						
4		· · · · · · · · · · · · · · · · · · ·			·	

Sources: (1) USEPA 2003, (2) WDOE 2002, (3) Bell 1986, (4) USEPA 2001, (5) Murray and McPhail 1988, (6) Brungs and Jones 1977

Murray and McPhail (1988) incubated five species of Pacific salmon, including coho, at five temperatures (2, 5, 8, 11, 14°C) to determine embryo survival at various temperatures. Coho embryos suffered increased mortality above 11°C although survival was still high. They concluded that the upper limit for normal coho embryo development is 14°C (Murray and McPhail 1988).

1.5 Freshwater Rearing and Growth

All of the freshwater rearing and growth temperature needs referenced in this section can be found in Table 8 through Table 10. Temperature affects metabolism, behavior, and survival of both juvenile fish as well as other aquatic organisms that may be food sources. In streams of the Northern California Coast, including the Klamath River, young Chinook, coho and steelhead may rear in freshwater from one to four years before migrating to the ocean.

In an exhaustive study of both laboratory and field studies of temperature effects on salmonids and related species, USEPA (1999a) concluded that temperatures of approximately 22-24°C limit salmonid distribution, i.e., they totally eliminate salmonids from a location. USEPA (1999a) also notes that changes in competitive interactions between fish species can lead to a transition in dominance from salmonids to other species at temperatures 2-4°C lower than the range of total elimination.

To protect salmon and trout during summer juvenile rearing the USEPA (2003) for Region 10 provided a single guidance metric designating 16°C as the 7-DADM

temperature that should not be exceeded in areas designated as "core" rearing locations. Core rearing areas are defined as areas with moderate to high densities of summertime salmonid juvenile rearing generally found in the mid- to upper portions of river basins. This criterion will protect juvenile salmonids from lethal temperatures, provide optimal to upper optimal conditions for juvenile growth depending on the time of year, avoid temperatures where salmonids are at a competitive disadvantage with other fish species, protect against increased disease rates caused by elevated temperatures, and provide temperatures which salmonids prefer according to scientific studies.

1.5.1 Steelhead Freshwater Rearing and Growth

Nielsen et al. (1994) studied thermally stratified pools and their use by juvenile steelhead in three California North Coast rivers including the Middle Fork Eel River, Redwood Creek at Redwood National Park, and Rancheria Creek, located in the Navarro River watershed. In detailed observations of juvenile steelhead behavior in and near thermally stratified pools in Rancheria Creek, Nielsen et al. (1994) noted behavioral changes including decreased foraging and increased aggressive behavior as pool temperature reached approximately 22°C. As pool temperature increased above 22°C, juveniles left the observation pools and moved into stratified pools where temperatures were lower.

Wurtsbaugh and Davis (1977, as cited by USEPA 2001) found that steelhead trout growth could be enhanced by temperature increases up to 16.5°C. Using a risk assessment approach which took into account "realistic food estimates", Sullivan et al. (2000) report temperatures of 13-17.0°C (MWAT), 14.5-21°C (MWMT), and 15.5-21°C (annual maximum) will ensure no more than a 10% reduction from maximum growth for steelhead. Reduction from maximum growth will be ≤20% for temperatures ranging from 10-19.0°C (MWAT), 10-24°C (MWMT), and 10.5-26°C (annual maximum).

A literature review was conducted by WDOE (2002) in which studies to determine the water temperature that would allow for maximum growth of steelhead trout were analyzed. These included laboratory studies conducted at constant and fluctuating temperatures. One of the studies was conducted using feeding rates comparable to those observed in natural creeks, although most of the laboratory studies were conducted under satiated feeding conditions. As a result of this review of laboratory studies conducted at constant temperatures, WDOE (2002) concludes that under satiated rations growth may be maximized at temperatures as high as 17.2-19°C. Results from laboratory studies using variable temperatures show maximum growth occurs at average daily temperatures between 15.5-18°C, and that under feeding rates similar to natural conditions at various times of the year maximum growth rates occurred at mean temperatures of 13.3°C (spring season), 15.2°C (fall season) and 16.2°C (summer season).

Table 8: Effects of Temperature in Considering Juvenile Steelhead Rearing and Growth

	e 8: Effects of Temperature in Con				'tn	
°C		REARI	NG AND GROWT	H		
26						21-26
25						Annual
	22-24 Temperature range which				21-24	maximum
24	totally eliminates salmonids from				MWMT	temp. which
	area, limiting their distribution (6)				which will	will ensure
23			left observation poo		ensure no	no more
23			with lower tempera	tures (2)	more than	than 20%
			ased foraging,	18-22	20%	reduction
22		increased ag	gressive behavior	Temperature	reduction	from max.
			(2)	range at	from max	growth (4)
				which	growth (4)	_
21				transition in		
				dominance		
				from		
20		17.10	17.2.10.5	salmonids to		
19		17-19	17.2-19 Growth	other species occurs (6)		
18		MWAT will ensure no	may be maximized at	occurs (0)		15.5-21
		more than	temperatures as			Annual
		20%	high as this			maximum
		reduction	under satiated		14.5-21	temperature
		from max.	feeding	15.5-18	MWMT	which will
17		growth (4)	conditions, lab	Average	which will	ensure no
		· · ·	studies at	daily	ensure no	more than
			constant	temperatures	more than	10% reduction
			temperature (5)	at which	10% reduction	from
	16.5 Growth enhanced by temp.		16 MWMT	maximum	from	maximum
	increases up to this temp. (3)	13-17	should not	growth occurs under	maximum	growth (4)
	16.2 Mean temp. at which max.	MWAT	exceed this value	satiated	growth (4)	growth (1)
16	growth occurred during the	range which	to be protective	feeding, lab	growth (1)	
	summer, lab studies using natural	will ensure	of core rearing	studies at		
	feeding conditions and varying	no more	locations (1)	varying		
	temps. (5)	than 10%	. ,	temps (5)		
	15.2 Mean temp. at which max.	reduction				
15	growth occurred during the fall, lab studies using natural feeding	from maximum				
	conditions and varying temps. (5)	growth (4)				10.5-15.5
	conditions and varying temps. (3)	510 will (7)		1		Annual
14						maximum
	13.3 Mean temp. at which max.				10-14.5	temperature
	growth occurred during the				MWMT	which will
13	spring, lab studies using natural	10-13			which will	ensure no
	feeding conditions and varying	MWAT will			ensure no	more than
	temps. (5)	ensure no			more than	20%
12	• • •	more than			20%	reduction
11		20%			reduction	from
		reduction			from	maximum
10		from			maximum	growth (4)
10		maximum			growth (4)	
	one (1) USEDA 2002 (2) Nielsen et e	growth (4)				

Sources: (1) USEPA 2003, (2) Nielsen et al. 1994, (3) Wurtsbaugh and Davis 1977, as cited by USEPA 2001, (4) Sullivan et al. 2000, (5) WDOE 2002, (6) USEPA 1999a

1.5.2 Chinook Freshwater Rearing and Growth

In a laboratory study, Brett (1952) demonstrated that juvenile Chinook salmon, acclimated to a temperature of 20°C, selectively aggregated in areas where the temperature was in the region of 12-13°C.

ODEQ (1995a), reviewed numerous studies and concluded for juvenile spring Chinook salmon rearing, positive growth takes place at temperatures between 4.5-19°C, and that optimum rearing production is between 10.0-15.6°C. However, as the extremes of this temperature range are reached growth reaches zero. Above and below these thresholds growth becomes negative as feeding ceases and respiration rates increase and/or decrease rapidly.

After synthesizing data from several sources USEPA (2001), came up with the same recommended optimum temperature zone for all Chinook salmon as ODEQ (1995a) of 10.0-15.6°C. While there is research suggesting that some Chinook stocks exhibit adequate rearing capabilities above 15.6°C, USEPA (2001) conclude that anything over this threshold significantly increases the risk of mortality from warm-water diseases.

In a laboratory study Marine and Cech (2004) studied the incremental effects of chronic exposure to three temperature regimes (13-16 °C, 17-20 °C, and 21-24 °C) on Chinook juveniles during rearing and smoltification. Their findings reflected that Chinook juveniles reared at the 17-20 °C and 21-24 °C temperature ranges experienced significantly decreased growth rates, impaired smoltification indices, and increased predation vulnerability compared with juveniles reared at 13-16 °C.

In a field study Chinook grew faster in a stream where temperatures peaked at 16°C compared to a stream where temperatures peaked at 20°C (ODFW 1992, as cited by WDOE 2002). WDOE (2002) reviewed literature on Chinook growth including laboratory studies conducted at a constant temperature, laboratory studies conducted at fluctuating temperatures, and field studies to evaluate the water temperature that would be protective of Chinook and allow for maximum growth. Most of the laboratory studies were conducted under satiated feeding conditions, although one of the studies was conducted using feeding rates more comparable to those observed in natural creeks. As a result of this review of laboratory studies conducted at constant temperatures, WDOE (2002) concludes that maximum growth is expected to occur with exposure to constant temperatures from 15.6-19°C. However, increased growth at temperatures above 15.6°C was inconsistently greater, and under natural rations the temperatures at which maximum growth occurs may decline by as much as 4.2°C. Recommendations based on the review of two laboratory studies conducted at fluctuating temperatures are that "...average temperatures below 19°C are necessary to support maximum growth rates in Chinook salmon, and that the average temperature that produces maximum growth rates likely lies between 15-18°C (median 16.5°C)".

Table 9: Effects of Temperature in Considering Juvenile Chinook Rearing and Growth

°C	ic). Effects of Tempera		ivenile Chinook Rearing RING AND GROWTH	g and Growth	
24 23 22 21	22-24 Temperature range which totally eliminates salmonids from area, limiting their distribution (7)			21-24 Decreased growth, impaired smoltification, increased predation compared to juveniles reared at 13-1 (6)	
20			18-22 Temperature		
19	19 Temperatures above this do not support maximum growth, lab studies at varying temperatures (3)	15.6-19 Maximum	range at which transition in dominance from salmonids to other species occurs (7)	17-20 Decreased growth, impaired smoltification, increased predation compared to juveniles	
18		growth expected according to lab		reared at 13-16 (6)	
17		studies conducted at constant temperature	16 Chinook grew		
16	15-18 Average temperature where maximum growth occurs, lab studies conducted at varying temperatures (3)	and satiated rations. Under natural feeding conditions maximum growth may occur at temperatures as much as 4.2C lower (3)	faster in a stream where temperatures peaked at 16 than when they peaked at 19C (3) 16 MWMT should not exceed this value to be protective of core rearing locations (2)	13-16 Increased growth, unimpaired smoltification, lower predation compared to juveniles reared at 21- 24, or 17-20 (6)	4.5-19 Temperature range at which
15					positive growth takes place (5)
14	10-15.6 Temperature range for optimal		10.15.6.0		
13	growth. Anything over this threshold		10-15.6 Optimal temperature range for	12-13 Juvenile Chinook acclimated to 20	
12	increases the risk of mortality from warm water disease (1)		rearing (5)	selectively aggregate to these water temperatures (4)	
11 10					
9			<u> </u>	<u> </u>	
8 7					
6					
5					
4					

Sources: (1) USEPA 2001, (2) USEPA 2003, (3) WDOE 2002, (4) Brett 1952, (5) ODEQ 1995a, (6) Marine and Cech 2004, (7) USEPA 1999a

1.5.3 Coho Freshwater Rearing and Growth

In a study of juvenile coho presence and absence in the Mattole watershed, Welsh et al. (2001) used logistic regression to determine that an MWAT greater than 16.8°C or a MWMT greater than 18.1°C may preclude the presence of juvenile coho salmon in the stream. The criterion correctly determined the presence or absence of juvenile coho in 18 of 21 streams. Welsh et al. (2001) also reported that juvenile coho were found in all streams with an MWAT less than 14.5°C, or a MWMT less than 16.3°C.

Sullivan et al. (2000) reviewed sub-lethal and acute temperature thresholds from a wide range of studies, incorporating information from laboratory-based research, field observations, and risk assessment approaches. Using a risk assessment approach based on "realistic food estimates" Sullivan et al (2000) suggest that MWATs ranging from 12.5-14.5°C for coho will result in no more than a 10% reduction from maximum growth, and that a range for the MWAT of 9-18.5°C will reduce growth no more than 20% from maximum. Sullivan et al. (2000) also calculated temperature ranges for MWMT (13-16.5°C) and the annual maximum temperature (13-17.5°C) that will result in no more than a 10% reduction in maximum growth. They further calculated ranges for MWMT (9-22.5°C) and the annual maximum temperature (9.5-23°C) that will result in no more than a 20% growth loss.

In an attempt to determine the water temperature that will allow for maximum growth of coho salmon, WDOE (2002) reviewed literature on laboratory studies conducted at a constant temperature and fluctuating temperatures, and field studies. The two laboratory studies reviewed were conducted under satiated feeding conditions. Shelbourn (1980, as cited by WDOE 2002) found that maximum growth occurred at a constant temperature of 17°C, while Everson (1973, as cited by WDOE 2002) tested fish at different temperatures and determined that coho had the greatest growth at the temperature test regime from 12.1-20.8°C (median 16.5°C). While the various field studies reviewed did not provide an estimate of the temperature best for maximum growth they did allow for WDOE (2002) to conclude that weekly average temperatures of 14-15°C were more beneficial to growth than lower temperature regimes, and daily maximum temperatures of 21-26°C were detrimental to growth.

Brett (1952) acclimated five different species of salmon to various temperatures ranging from 5-24°C and found that coho salmon showed the greatest preference for temperatures between 12-14°C. It was also determined that coho showed a general avoidance of temperatures above 15°C even in fish who were acclimated to temperatures as high as 24°C.

Konecki et al. (1995a) raised two groups of juvenile coho salmon under identical regimes to test the hypothesis that the group from a stream with lower and less variable temperature would have a lower and less variable preferred temperature than the group from a stream with warmer and more variable temperatures. Results reflected that the two groups tended to differ in their preferred temperature range as predicted above, but the differences were slight. Konecki et al. (1995a) concluded that the temperature preference of juvenile coho salmon in their study was 10-12°C.

Table 10: Effects of Temperature in Considering Juvenile Coho Rearing and Growth

°C	ble 10: Effects of Temperature in Considering Juvenile Coho Rearing and Growth REARING AND GROWTH				
26		KE/	KING MND GRO	VV 111	
25					
24			22-24 Tempe	rature range which totally	
23	21-26 Daily maximum temperatures in this range are detrimental to growth, according to			ids from an area, limiting their	
22			d	istribution (9)	
22	field studies (
21	1010 334003			18-22 Temperature range at	17.5-23 Annual maximum temperature will ensure no
20				which transition in	more than 20%
19				dominance from salmonids	reduction from
18	18.1 MWMT above this may preclude the presence of juvenile coho in steams (5)		16.5-22.5 MWMT will ensure no more than 20%	to other species occurs (9)	maximum growth (2)
			reduction from	17 Maximum growth at this	
15			maximum	constant temperature, at	
17			growth (2)	satiated rations in a lab study	
				(6)	
	12.1-20.8		16.8 MWAT above this may		
			preclude the presence of		
			juvenile coho in streams (5)		
1.0	14.5-18.5 MWAT will	Greatest growth		16.3 Juveniles found in all	
16	ensure no more than 20%	occurs in this		streams with MWMT less	13.5 17.5
	reduction from maximum	temperature		than this value (5) 16 MWMT not exceed this	Annual
	growth (2)	range under		value to be protective of core	maximum
		satiated		rearing locations (1)	temperature
		conditions, lab		>15 Juveniles show	will ensure no
		study (7)		avoidance, even those	more than 10%
15			13-16.5 MWMT	acclimated to 24C (4)	reduction from maximum
			will ensure no	14-15 Weekly average	maximum growth (2)
			more than 10%	temperatures in this range	giowiii (2)
			reduction from	are more beneficial than	
1.	1451		maximum	lower temperatures (3)	
14	14.5 Juvenile coho found in all streams with MWAT		growth (2)		
	less than this value (5)				
	12.5-14.5 MWAT will			12-14 Preferred temperature	
	ensure no more than 10%			range (4)	
13	reduction from maximum			1	9.5-13.5
	growth (2)		0.1234000		Annual
12			9-13 MWMT will ensure no	10.12 Par Com. 1	maximum
11	9-12.5 MWAT will ensure		more than 20%	10-12 Preferred temperature	temperature
10	no more than 20%		reduction from	range (8)	will ensure no
	reduction from maximum		maximum		more than 20%
9	growth (2)		growth (2)		reduction from
					max. growth
					(2)

Sources: (1) USEPA 2003, (2) Sullivan et al. 2000, (3) WDOE 2002, (4) Brett 1952, (5) Welsh et al. 2001, (6) Shelbourn 1980, as cited by WDOE 2002, (7) Everson 1973, as cited by WDOE 2002, (8) Konecki et al. 1995a, (9) USEPA 1999a

1.6 Lethality

All of the lethal temperatures referenced in this section can be found in Table 11. WDOE (2002) reviewed literature on three types of studies (constant exposure temperature studies, fluctuating temperature lethality studies, and field studies) and used this information to calculate the MWMT that, if exceeded, may result in adult and juvenile salmonid mortality. The resultant MWMTs for these various types of studies are as follows: constant exposure studies 22.64°C, fluctuating lethality studies 23.05°C, and field studies 22.18°C.

1.6.1 Steelhead Lethality

Coutant (1970, as cited by USEPA 1999a) found that Columbia River steelhead, which were acclimated to a river temperature of 19°C, had a lethal threshold of 21°C. Bell (1986) reviewed various studies and states that the lethal threshold for steelhead is 23.9°C. According to the California Department of Fish and Game (2001, p.419), temperatures of 21.1°C have been reported as being lethal to adults.

1.6.2 Chinook Lethality

In a laboratory study, Brett (1952) acclimated five different species of juvenile salmon to various temperatures ranging from 5-24°C. At temperatures of 24°C and below there was 100% survival of fish during the one-week duration of the experiment. Brett (1952) concluded that the lethal temperature (temperature where survival becomes less than 100%) was between 24.0 and 24.5°C, and the ultimate upper lethal temperature was 25.1°C (temperature at which 50% of the population is dead after infinite exposure). A review of numerous studies led Bell (1986) to conclude that the upper lethal temperature for Chinook is 25°C. Myrick and Cech (2001) reviewed literature on studies from the Central Valley and found data to suggest that the chronic (exposure >7 days) upper lethal limit for juvenile Chinook is approximately 25°C.

1.6.3 Coho Lethality

In a review of various literature sources, Bell (1986) found that the upper lethal temperature for coho is 25.6°C. Brett (1952) concluded that the ultimate upper lethal temperature of juvenile coho salmon was 25.0°C (temperature at which 50% of the population is dead after infinite exposure). Thomas et al. (1986) conducted a study to determine the mortality of coho subjected to fluctuating temperatures. It was determined that the LT50 (the temperature at which 50% of the population will die) for fish acclimated to a 10-13°C cycle was 26°C for presmolts (age-2 fish), and 28°C for age-0 fish.

Table 11: Effects of Temperature in Considering Lethality and Salmonids

°C	Steelhead	Chinook	Coho	All Salmonids
28			28 LT50 ¹ for age 0-fish acclimated to a 10-13C cycle (6)	
27				
26			26 LT50 ¹ for presmolts (age 2-fish) acclimated to a 10- 13C cycle (6)	
25		25.1 Upper lethal temp. at which 50% of the population would die after infinite exposure, juvenile Chinook acclimated to temperatures from 5-24C (4)	25.6 Upper lethal threshold (3)	
25		25 Upper lethal threshold (3) 25 Chronic (exposure >7	25 Upper lethal temp. at which 50% of the population would die after	
		days) upper lethal limit for juvenile Chinook (5).	infinite exposure, juvenile coho acclimated to temps. from 5-24C (4)	
24		24-24.5 Survival becomes less than 100% for juvenile Chinook acclimated to temperatures from 5-24C (4)		
23	23.9 Upper lethal threshold for steelhead (3)			23.05 do not exceed this value to prevent adult and juvenile mortality, data from fluctuating temp. studies (1)
22				22.64 do not exceed this value to prevent adult and juvenile mortality, data from constant exposure studies (1) 22.18 do not exceed this value to prevent adult and juvenile mortality, data from field studies (1)
21	21.1 Temperature lethal to adults (7) 21 Lethal threshold for steelhead acclimated to 19C (2)			

¹ Maximum temperature in the cycle at which 50% mortality occurred

Sources: (1) WDOE 2002, (2) Coutant 1970, as cited by USEPA 1999a, (3) Bell 1986, (4) Brett 1952, (5) Myrick and Cech 2001, (6) Thomas et al. 1986, (7) California Department of Fish and Game (CDFG) 2001

1.7 Disease

All of the effects of temperatures on disease risk in salmonids referenced in this section can be found in Table 12. WDOE (2002) reviewed studies of disease outbreak in salmonids and estimates that an MWMT of less than or equal to 14.38°C (midpoint of 12.58-16.18 range) will virtually prevent warm water disease effects. To avoid serious

19

Table 12: Effects of Temperature in Considering Disease and Salmonids

°C		ch	Considering Disease and Ceratomyxos		Columnaris	Disease (general)	
26	1	CII	Ceratomyxos	015	Columnaris	Disease (general)	
25							
24	>24 Lifecycle takes less than 4 days (5)	21-26.7					
23	21-23.9 Life cycle takes as few as 3-4 days (5)	Optimum temp. range for Ich, compilation of temps. from three references (3,4,5)	23.3 Juvenile coho salmon and rainbow trout time from exposure to death is 12.5 and 14 days respectively (9)		23.3 Juvenile spring Chinook mortality was 92%, and time from exposure to death was 2.3 days (13)		
22					22.2 Mortality is 100% in juvenile sockeye exposed to <i>C. columnaris</i> (10)		
21					>21.1 Temperatures at this level are associated with a 28-74% infection rate in Chinook (11)		
20	18.3-21.2 Serious outbreaks of Ich occur (4)	20 Lifecycle takes 1 week (6)	20.5 Mortality is 84% in juvenile coho exposed to <i>C. shasta</i> (9).	6.7-23.3 Juvenile rainbow trout have little or no ability to overcome infection,	20.5 Mortality in juvenile steelhead and coho from Columnaris was 100%, and 70% in juvenile spring Chinook (13) 20.5 In juvenile steelhead and coho time from exposure to death was 1.6-1.7 days (13) 20 Average water temperature at which low virulence strains show signs of outbreak (3, 12)	>20.88 MWMTs over this value can result in severe infections and catastrophic outbreaks (1)	18-20 Temperature range which is associated with a high risk of disease in rearing juveniles
19				and mortality			and migrating
18				varied			adults (2)
17				from 75- 86% (9)	17.8 Mortality rates were 52, 92, and 99% for juvenile spring Chinook, steelhead and coho respectively (13)	17.38 MWMT should not be exceeded to avoid serious rates of infection and mortality (1)	14-17 Temperature
16					16.1 Mortality is 30% in juvenile sockeye exposed to <i>C. columnaris</i> (10)		range which is associated
15	in salmonid especially (15.5 Lifecycle	ed with outbreaks I fingerlings, Chinook (3) e of Ich takes 2 ks (5)	15 Mortality is 22% in juvenile coho exposed to <i>C. shasta</i> (9).		15.6 Average water temperature at which low virulence strains show signs of outbreak (3) 15 Mortality was 31, 56, and 51% for juvenile spring Chinook, steelhead, and coho respectively (13)		with an elevated risk of disease in rearing juveniles and
14						14.38 MWMT will virtually prevent all warm water disease (1)	migrating adults (2)

Table 12 (continued): Effects of Temperature in Considering Disease and Salmonids

°C	Ich	Ceratom	yxosis	Columnaris	Disease (general)
13 12 11 10	10 Lifecycle takes more than 5 weeks (5)	10-11 <i>C. shasta</i> appears to be come infective (4) <10 Steelhead show evidence of <i>C. shasta</i> in ~38 days (8) 9.4 Juvenile coho time from exposure to death is 146 days, mortality is 2% (9)	6.7-23.3 Juvenile rainbow trout have little or no ability to overcome infection, and mortality varied from 75-86% (9)	12.8 After 7 days of infection mortality is 60-100% (majority of tests 100%) (12) 12.2 Mortality was 4-20% in juvenile spring Chinook, steelhead, and coho respectively. Time from exposure to death ranged from 7.6-12.2 days (13).	12-13 Temperature range which minimizes the risk of disease in rearing juveniles and migrating adults (2)
7	7 Lifecycle takes 20 days (6) <7 Lifecycle takes more than 5 weeks (7)				
5 4 3			6.7 Juvenile rainbow trout time from exposure to death is 155 days (9) 3.9-6.7 No mortality in Juvenile coho exposed to <i>C. shasta</i> (9)	3.9-9.4 No mortality in spring Chinook, steelhead, or coho from Columnaris (13)	

Sources: (1) WDOE 2002, (2) USEPA 2003, (3) Bell 1986, (4) CDWR 1988, (5) Piper et al 1982, (6) Nigrelli et al. 1976, as cited by Dickerson et al. 1995, (7) Durborow et al. 1998, (8) Leitritz and Lewis, 1976, (9) Udey et al. 1975, (10) Ordal and Rucker 1944, as cited by Pacha et al. 1970, (11) USEPA 1999a, (12) Pacha et al. 1970, (13) Holt et al. 1975

rates of infection and mortality the MWMT should not exceed 17.38°C (midpoint of 15.58-19.18 range), and that severe infections and catastrophic outbreaks become a serious concern when the MWMTs exceed 20.88°C (midpoint of 18.58-23.18 range).

In a summary of temperature considerations, USEPA (2003) states that disease risks for juvenile rearing and adult migration are minimized at temperatures from 12-13°C, elevated from 14-17°C, and high at temperatures from 18-20°C.

Acknowledging that there are many diseases that affect salmonids, the following discussion will focus on three which are common in the Klamath Basin: Ichthyophthiriasis (Ich), Ceratomyxosis, and Columnaris. *Ichthyophthirius multifiliis* is a protozoan parasite that causes the disease known as Ichthyophthiriasis (Ich). The disease ceratomyxosis is caused by a parasite, *Ceratomyxa shasta* (*C. shasta*). Columnaris disease is a bacterial infection caused by *Flavobacterium columnare* (synomyms: *Bacillus columnaris*, *Chondrococcus columnaris*, *Cytophaga columnaris*, *Flexibacter columnaris*).

1.7.1 Ichthyophthiriasis (Ich)

Nigrelli et al. (1976, as cited by Dickerson et al. 1995) proposed that there are physiological races of Ich, which are related to the temperature tolerance of the host fishes. Thus, there are races of Ich that infect cold-water (7.2-10.6°C) fishes such as salmon, and others that infect warm-water (12.8-16.1°C) tropical fishes. Bell (1986) discusses Ich and states that at water temperatures above 15.6°C, this disease often breaks out in salmon fingerlings, especially Chinook. CDWR (1988) states that serious outbreaks of Ich occur at temperatures from 18.3-21.2°C.

Numerous studies and reviews have been conducted on the optimal temperature for Ich. Piper et al. (1982, p.316.) wrote that optimal temperatures range from 21-23.9°C. CDWR (1988) stated the optimum temperature for Ich is in the range of 25 to 26.7°C, while Bell (1986) states optimum temperatures are noted from 21.2-26.7°C.

Temperature is an important factor in the persistence of Ich infections in salmonids. The growth period varies from 1 week at 20 °C to 20 days at 7 °C (Nigrelli et al. 1976, as cited by Dickerson et al. 1995). Piper et al. (1982, p.316) state that at optimal temperatures of 21-23.9 °C, the life cycle may take as few as 3-4 days. The cycle requires 2 weeks at 15.5 °C, and more than 5 weeks at 10 °C (Piper et al. 1982, p.316). Durborow et al. (1998) note that to complete its lifecycle, Ich requires from less than 4 days at temperatures higher than 24 °C, to more than 5 weeks at temperatures lower than 7 °C. Although studies report varying lengths of time for Ich to complete its lifecycle at similar temperatures, it is clear that the speed at which Ich develops increases as temperatures increase.

1.7.2 Ceratomyxosis

In reviewing the literature on Ceratomyxosis (a disease caused by the parasite, C. shasta), it is clear that as water temperatures increase, the intensity of the disease increases, and the incubation period decreases (CDWR 1988, Letritz and Lewis, Udey et al. 1975). At

water temperatures greater than 10°C, steelhead will show evidence of Ceratomyxosis in approximately 38 days (Leitritz and Lewis 1976, p.154). In a study of juvenile coho salmon by Udey et al. (1975), time from exposure to death was more than 90% temperature dependent, and increased from 12.5 days at 23.3°C, to 146 days at 9.4°C. These results show the accelerating effect of higher temperatures on the progress of the disease. The time from exposure to death of juvenile rainbow trout was nearly 97% temperature dependent, increasing from 14 days at 23.3°C to 155 days at 6.7°C (Udey et al. 1975).

C. shasta appears to become infective at temperatures around 10-11°C (CDWR 1988). According to Leitritz and Lewis (1976, p.154), steelhead from the Klamath River are quite susceptible to *C. shasta* infections and suffer severe losses when exposed.

Udey et al. (1975) conducted a study to determine the relation of water temperature to Ceratomyxosis in juvenile rainbow trout and coho salmon. Rainbow trout from the Roaring River Hatchery, and coho from Fall Creek Salmon Hatchery (both in Oregon) were used in this experiment. Groups of 25 fish exposed to *C. shasta* were transferred to 12.2°C water, and then were tempered to one of eight experimental temperatures from 3.9 to 23.3°C (2.8°C increments).

In the juvenile coho salmon experiment, Udey et al. (1975) found that percent mortality increased progressively from 2% at 9.4°C to 22% at 15.0°C and 84% at 20.5°C. No deaths occurred in coho salmon maintained at 3.9 and 6.7°C, indicating that ceratomyxosis in coho can be suppressed by water temperatures of 6.7°C or below (Udey et al. 1975).

Tests conducted by Udey et al. (1975) on rainbow trout juveniles indicate that once infection is initiated, juvenile rainbow trout have little or no ability to overcome *C. shasta* infections at water temperatures between 6.7 and 23.3°C. Fatal infections varied from 75-86% at temperatures ranging from 6.7 to 15.0°C (Udey et al. 1975). Mortality in trout held at 20.5 and 23.3°C were lower (72% and 52% respectively) due to losses from *Flexibacter columnaris*, which occurred well before the onset of deaths caused by *C. shasta*, in spite of efforts to control it with terramycin (Udey et al. 1975). The results from Udey et al. (1975) also reflected no deaths occurred in juvenile trout held at 3.9°C.

1.7.3 Columnaris

The importance of temperature on infections of Columnaris has been demonstrated in numerous laboratory studies. Ordal and Rucker (1944, as cited by Pacha et al. 1970) exposed juvenile sockeye salmon to *C. columnaris* and studied the effect of temperature on the disease. In these studies, the overall mortality ranged from 30% in fish held at 16.1°C to 100% in those held at 22.2°C (Ordal and Rucker 1944, as cited by Pacha et al. 1970). USEPA (1999a) cites studies that conducted surveys of Columnaris infection frequency on Chinook in the Snake River in July and early August of 1955-1957, which revealed 28-75% of fish infected when water temperature was >21.1°C.

Low virulence strains of Columnaris show signs of outbreak when average water temperatures are over 20°C (Bell 1986, Pacha et al. 1970). Bell (1986) states that

outbreaks of high virulence strains occur when average water temperatures reach 15.6°C, and Pacha et al. (1970) found mortalities of 60-100% (majority of tests 100%) occur at temperatures of 12.8°C after 7 days of infection. With regard to strains of higher virulence, while these strains are capable of beginning infection and producing disease at water temperatures as low as 12.8°C, the disease process becomes progressively slower as the water temperature is lowered (Pacha et al. 1970).

Holt et al. (1975) performed a study on the relation of water temperature to Columnaris in juvenile steelhead trout and juvenile coho and spring Chinook salmon. Tests were performed on groups of 25-35 fish at eight temperatures ranging from 3.9°C to 23.3°C (2.8°C increments). At 20.5°C mortality was 100% in juvenile steelhead trout and coho salmon, 70% in juvenile spring Chinook salmon, and at temperatures 23.3°C juvenile spring Chinook mortality was 92% (Holt et al. 1975). Mortality rates were 52, 92, and 99% at 17.8°C for juvenile spring Chinook, steelhead trout, and coho salmon respectively, and mortality dropped to 31, 56, and 51% at 15.0°C (Holt et al. 1975). At 12.2°C mortality varied from 4 to 20% among juveniles of the three species, and at temperatures of 9.4°C and below, no deaths due to the experimental infection with *F. columnaris* occurred (Holt et al. 1975). Holt et al. (1975) state that these results indicate that under the conditions of these experiments Columnaris disease was completely suppressed by water temperatures of 9.4°C or below.

In general, data from laboratory studies indicates that as water temperatures increase, the time to death decreases (Pacha et al. 1970). With juvenile steelhead trout and juvenile coho and spring Chinook salmon as the temperature increased above 12.2°C, the disease process was progressively accelerated, resulting in a minimum time to death at 20.5 or 23.3°C and a maximum at 12.2°C (Holt et al. 1975). In these juvenile salmonids Holt et al. (1975) found the mean time to death decreased from 7.6-12.2 days at 12.2°C to 1.6-1.7 days at 20.5°C for juvenile coho and steelhead, and 2.3 days at 23.3°C for juvenile spring Chinook (Holt et al. 1975).

1.8 TMDL Temperature Thresholds

Currently there are no numeric temperature standards in the *Water Quality Control Plan* for the North Coast Region (Basin Plan). Thus, information from this literature review will be utilized by Regional Water Board staff to selected chronic and acute temperature thresholds for evaluation of stream temperatures in TMDLs. Chronic temperature thresholds (MWMTs) were selected from the USEPA document *EPA Region 10 Guidance For Pacific Northwest State and Tribal Temperature Water Quality Standards* (2003), and are presented in Table 13. The Region 10 guidance is the product of a three-year interagency effort, and has been reviewed by both independent science review panels and the public. Acute lethal temperature thresholds were selected based upon best professional judgment of the literature, and are presented in Table 14.

Table 13: Life Stage Temperature Thresholds

Life Stage	MWMT (°C)
Adult Migration	20
Adult Migration plus Non-Core ¹ Juvenile Rearing	18
Core ² Juvenile Rearing	16
Spawning, Egg Incubation, and Fry Emergence	13

¹ Non-Core is defined as moderate to low density salmon and trout rearing usually occurring in the mid or lower part of the basin (moderate and low not defined).

Table 14: Lethal Temperature Thresholds

Lethal Threshold ¹ (°C)					
Life Stage Steelhead Chinook Coho					
Adult Migration and Holding	24	25	25		
Juvenile Growth and Rearing	24	25	25		
Spawning, Egg Incubation, and Fry Emergence	20	20	20		

¹ The lethal thresholds selected in this table are generally for chronic exposure (greater than seven days). Although salmonids may survive brief periods at these temperatures, they are good benchmarks from the literature for lethal conditions.

In some cases it may be necessary to calculate MWATs for a given waterbody, and compare these to MWAT thresholds. USEPA (2003) states that for many rivers in the Pacific Northwest the MWMT is about 3°C higher than the MWAT (USEPA 2003, as cited by Dunham et al. 2001and Chapman 2002). Rather than list MWAT thresholds in this document using the 3°C difference suggested above, the Regional Water Board will consider stream temperatures within each individual TMDL waterbody. Thus the Regional Water Board will calculate both MWMTs and MWATs for the waterbody, and characterize the actual difference between these temperature metrics for the watershed using an approach similar to that used in Sullivan et al. (2000). Once this relationship is understood, MWAT thresholds for each life stage can be identified and compared to the watershed MWATs.

The freshwater temperature thresholds presented in this section are applicable during the season or time of year when the life stage of each species is present. Periodicity information is not discussed in this document and will be presented in each individual TMDL staff report. Where life history, timing, and/or species needs overlap, the lowest of each temperature metric applies.

² Core is defines as areas of high density rearing (high is not specifically defined). Source: USEPA 2003

CHAPTER 2. DISSOLVED OXYGEN and TOTAL DISSOLVED GAS

2.1 Introduction

Adequate concentrations of dissolved oxygen in fresh water streams are critical for the survival of salmonids. Fish have evolved very efficient physiological mechanisms for obtaining and using oxygen in the water to oxygenate the blood and meet their metabolic demands (WDOE 2002). Reduced levels of dissolved oxygen can impact growth and development of different life stages of salmon, including eggs, alevins, and fry, as well as the swimming, feeding and reproductive ability of juveniles and adults. Such impacts can affect fitness and survival by altering embryo incubation periods, decreasing the size of fry, increasing the likelihood of predation, and decreasing feeding activity. Under extreme conditions, low dissolved oxygen concentrations can be lethal to salmonids. High levels of total dissolved gas concentrations (TDG), including dissolved oxygen, can result in gas bubble disease and death for salmonids.

Literature reviewed for this analysis included EPA guidance, other states' standards, reports that compiled and summarized existing scientific information, and numerous laboratory studies. When possible, species-specific requirements were summarized for the following life stages: migrating adults, incubation and emergence, and freshwater rearing and growth. The following information applies to salmonids in general, with specific references to coho, Chinook, steelhead, and other species of salmonids as appropriate.

2.2 Effects of Low Dissolved Oxygen Concentrations on Salmonids

2.2.1 Adult Migration

Reduced concentrations of dissolved oxygen can negatively affect the swimming performance of migrating salmonids (Bjornn and Reiser 1991). The upstream migration by adult salmonids is typically a stressful endeavor. Sustained swimming over long distances requires high expenditures of energy and therefore requires adequate levels of dissolved oxygen. Migrating adult Chinook salmon in the San Joaquin River exhibited an avoidance response when dissolved oxygen was below 4.2 mg/L, and most Chinook waited to migrate until dissolved oxygen levels were at 5 mg/L or higher (Hallock et al. 1970).

2.2.2 Incubation/Emergence

Low levels of dissolved oxygen can be directly lethal to salmonids, and can also have sublethal effects such as changing the rate of embryological development, the time to hatching, and size of emerging fry (Spence et al. 1996). The embryonic and larval stages of salmonid development are especially susceptible to low dissolved oxygen levels as their ability to extract oxygen is not fully developed and their relative immobility inhibits their ability to migrate to more favorable conditions. The dissolved oxygen requirements for successful incubation of embryos and emergence of fry is tied to intragravel dissolved oxygen levels. Intragravel dissolved oxygen is typically a function of many chemical, physical, and hydrological variables, including: the dissolved oxygen concentration of the

overlying stream water, water temperature, substrate size and porosity, biochemical oxygen demand of the intragravel water, sediment oxygen demand, the gradient and velocity of the stream, channel configuration, and depth of water. As a result the dissolved oxygen concentration within the gravels can be depleted causing problems for salmonid embryos and larvae, even when overlying surface water oxygen levels are suitable (USEPA 1986a).

Studies note that water column dissolved oxygen concentrations are typically estimated to be reduced by 1-3 mg/L as water is transmitted to redds containing developing eggs and larvae (WDOE 2002). USEPA (1986a) concluded that dissolved oxygen levels within the gravels should be considered to be at least 3 mg/L lower than concentrations in the overlying water. ODEQ (1995b) expect the loss of an average of 3 mg/L dissolved oxygen from surface water to the gravels.

2.2.3 Incubation Mortality

Phillips and Campbell (1961, as cited by Bjornn and Reiser 1991) concluded that intragravel dissolved oxygen must average 8 mg/L for embryos and alevins to survive well. After reviewing numerous studies Davis (1975) states that a dissolved oxygen concentration of 9.75 mg/L is fully protective of larvae and mature eggs, while at 8 mg/L the average member of the incubating population will exhibit symptoms of oxygen distress, and at 6.5 mg/L a large portion of the incubating eggs may be affected. Bjornn and Reiser (1991) reviewed numerous references and recommend that dissolved oxygen should drop no lower than 5 mg/L, and should be at or near saturation for successful incubation.

In a review of several laboratory studies, ODEQ (1995b) concluded that at near optimum (10°C) constant temperatures acute mortality to salmonid embryos occurs at relatively low concentrations of dissolved oxygen, near or below 3 mg/L. Field studies reviewed by ODEQ (1995b) demonstrate that embryo survival is low when the dissolved oxygen content in the gravels drops near or below 5 mg/L, and survival is greater at 8 mg/L.

Silver et al. (1963) performed a study with Chinook salmon and steelhead trout, rearing eggs at various constant dissolved oxygen concentrations and water velocities. They found that steelhead embryos held at 9.5°C and Chinook salmon embryos held at 11°C experienced complete mortality at dissolved oxygen concentrations of 1.6 mg/L. Survival of a large percentage of embryos reared at oxygen levels as low as 2.5 mg/L appeared to be possible by reduction of respiration rates and consequent reduction of growth and development rates.

In a field study Cobel (1961) found that the survival of steelhead embryos was correlated to intragravel dissolved oxygen in the redds, with higher survival at higher levels of dissolved oxygen. At 9.25 mg/L survival was 62%, but survival was only 16% at 2.6 mg/L. A laboratory study by Eddy (1971) found that Chinook salmon survival at 10.4 mg/L (13.5 °C) was approximately 67%, however at dissolved oxygen levels of 7.3 mg/L (13.5 °C) survival dropped to 49-57.6%. At temperatures more suitable for Chinook incubation (10.5 °C) Eddy (1971) found the percent survival remained high (over 90%) at

dissolved oxygen levels from 11 mg/L to 3.5 mg/L; however, as dissolved oxygen levels decreased, the number of days to hatching increased and the mean dry weight of the fry decreased substantially. WDOE (2002) also points out that the studies above did not consider the act of emerging through the redds, and the metabolic requirements to emerge would be expected to be substantial. Therefore, it is likely that higher oxygen levels may be needed to fully protect hatching and emergence, than to just support hatching alone.

2.2.4 Incubation Growth

Embryos can survive when dissolved oxygen is below saturation (and above a critical level), but development typically deviates from normal (Bjornn and Reiser 1991). Embryos were found to be smaller than normal, and hatching either delayed or premature, when dissolved oxygen was below saturation throughout development (Doudoroff and Warren 1965, as cited by Bjornn and Reiser 1991).

Garside (1966) found the number of days it took for rainbow trout to go from fertilization to hatching increased as dissolved oxygen concentrations and water temperature decreased. In this study, rainbow trout were incubated at temperatures between 2.5 - 17.5°C and dissolved oxygen levels from 2.5 - 11.3 mg/L. At 10°C and 7.5°C the total time for incubation was delayed 6 and 9 days respectively at dissolved oxygen levels of 2.5 mg/L versus embryos incubated at approximately 10.5 mg/L.

Silver et al. (1963) found that hatching of steelhead trout held at 9.5°C was delayed 5 to 8 days at dissolved oxygen concentrations averaging 2.6 mg/L versus embryos reared at 11.2 mg/L. A smaller delay of hatching was observed at oxygen levels of 4.2 and 5.7 mg/L, although none was apparent at 7.9 mg/L. For Chinook salmon held at 11°C, Silver et al. observed that embryos reared at oxygen levels lower than 11 mg/L experienced a delay in hatching, with the most significant delay in those reared at dissolved oxygen levels of 2.5 mg/L (6 to 9 days). The size of both Chinook and steelhead embryos increased with increases in dissolved oxygen up to 11.2 mg/L. External examination of embryos revealed abnormal structural development in Chinook salmon tested at dissolved oxygen concentrations of 1.6 mg/L, and abnormalities in steelhead trout at concentrations of 1.6 and 2.6 mg/L. The survival of Chinook salmon after hatching was only depressed at the 2.5 mg/L level, the lowest level at which hatching occurred, with lower mortalities occurring at higher velocities. Post hatching survival of steelhead trout could not be determined due to numerous confounding factors.

Shumway et al. (1964) conducted a laboratory study to determine the influence of oxygen concentration and water movement on the growth of steelhead trout and coho salmon embryos. The experiments were conducted at a temperature of 10°C and oxygen levels generally ranging from 2.5 - 11.5 mg/L and flows from 3 to 750 cm/hour. It was concluded that the median time to hatching decreased and size of fry increased as dissolved oxygen levels increased. For example, steelhead trout embryos reared at 2.9 mg/L hatched in approximately 41 days and had a wet weight of 17 mg, while embryos reared at 11.9 mg/L hatched in 36 days and weighed 32.3 mg. The authors found that a reduction of either the oxygen concentration or the water velocity will reduce the size of

fry and increase the incubation period, although the affect of various water velocities tested was less than the effect of the different dissolved oxygen concentrations tested.

WDOE (2002) reviewed various references and found that at favorable incubation temperatures a mean oxygen concentration of 10.5 mg/L will result in a 2% reduction in growth. At other oxygen concentrations, growth is reduced as follows: 8% reduction at oxygen levels of 9 mg/L, 10% reduction at 7 mg/L, and a 25% reduction at 6 mg/L.

2.2.5 Incubation Avoidance/Preference

Alevin showed a strong preference for oxygen concentrations of 8 - 10 mg/L and moved through the gravel medium to these concentrations, avoiding concentrations from 4 - 6 mg/L (WDOE 2002).

2.2.6 Emergence Mortality

"The hatching time, size, and growth rate of developing embryos is proportional to the dissolved oxygen concentrations up to 8 mg/L or greater. The ability of fry to survive their natural environment may be related to the size of fry at hatch (ODEQ 1995b)." McMahon (1983) recommends dissolved oxygen levels be ≥ 8 mg/L for high survival and emergence of fry. In a review of controlled field and lab studies on emergence, WDOE (2002) states that average intragravel oxygen concentrations of 6 - 6.5 mg/L and lower can cause stress and mortality in developing embryos and alevin. It is also noted that field studies on emergence consistently cite intragravel oxygen concentrations of 8 mg/L or greater as being associated with or necessary for superior health and survival, oxygen concentrations below 6 - 7 mg/L result in a 50% reduction in survival through emergence, and oxygen concentrations below 5 mg/L result in negligible survival. According to various laboratory studies, the threshold for complete mortality of emerging salmonids is noted to occur between 2 - 2.5 mg/L (WDOE 2002).

After reviewing numerous literature sources, the USEPA (1986a) concluded that the embryonic and larval stages of salmonid development will experience no impairment when water column dissolved oxygen concentrations are 11 mg/L. This translates into an intragravel dissolved oxygen concentration of 8 mg/L (USEPA assumes a 3 mg/L loss between the surface water and gravels). Table 15 from the USEPA (1986a) lists the water column and intragravel dissolved oxygen concentrations associated with various health effects. These health affects range from no production impairment to acute mortality.

Table 15: Dissolved oxygen concentrations and their effects salmonid embryo and larval stages

Level of Effect	Water Column DO (mg/L)	Intragravel DO (mg/L)
No Production Impairment	11	8*
Slight Production Impairment	9	6*
Moderate Production Impairment	8	5*
Severe Production Impairment	7	4*
Limit to Avoid Acute Mortality	6	3*

^{*} A 3 mg/L loss is assumed between the water column dissolved oxygen levels and those intragravel. Source: USEPA 1986a

2.2.7 Freshwater Rearing and Growth

2.2.7.1 Swimming and Activity

Salmonids are strong active swimmers requiring highly oxygenated waters (Spence 1996), and this is true during the rearing period when the fish are feeding, growing, and avoiding predation. Salmonids may be able to survive when dissolved oxygen concentrations are low (<5 mg/L), but growth, food conversion efficiency, and swimming performance will be adversely affected (Bjornn and Reiser 1991). Davis (1975) reviewed numerous studies and reported no impairment to rearing salmonids if dissolved oxygen concentrations averaged 9 mg/L, while at oxygen levels of 6.5 mg/L "the average member of the community will exhibit symptoms of oxygen distress", and at 4 mg/L a large portion of salmonids may be affected. Dahlberg et al. (1968) state that at temperatures near 20°C any considerable decrease in the oxygen concentration below 9 mg/L (the air saturation level) resulted in some reduction of the final swimming speed. They found that between dissolved oxygen concentrations of 7 to 2 mg/L the swimming speed of coho declined markedly with the decrease in dissolved oxygen concentration.

In a laboratory study, Davis et al. (1963) reported that the maximum sustainable swimming speeds of wild juvenile coho salmon were reduced when dissolved oxygen dropped below saturation at water temperatures of 10, 15, and 20°C. Air-saturation values for these dissolved oxygen concentrations were cited as 11.3, 10.2, and 9.2 mg/L respectively. They found that the maximum sustained swimming speeds (based on first and second swimming failures at all temperatures) were reduced by 3.2 - 6.4%, 5.9 - 10.1%, 9.9 - 13.9%, 16.7 - 21.2%, and 26.6 - 33.8% at dissolved oxygen concentrations of 7, 6, 5, 4, and 3 mg/L respectively. The authors also conducted tests on juvenile Chinook salmon and found that the percent reductions from maximum swimming speed at temperatures ranging from 11 to 15°C were greater than those for juvenile coho. At the dissolved oxygen concentrations listed above swimming speeds were decreased by 10%, 14%, 20%, 27%, and 38% respectively.

WDOE (2002) reviewed various data and concluded that swimming fitness of salmonids is maximized when the daily minimum dissolved oxygen levels are above 8 - 9 mg/L. Jones et al. (1971, as cited by USEPA 1986a) found the swimming speed of rainbow trout was decreased 30% from maximum at dissolved oxygen concentrations of 5.1 mg/L and 14°C. At oxygen levels of 3.8 mg/L and a temperature of 22°C, they found a 43% reduction in the maximum swimming speed.

2.2.7.2 Growth

In a review of constant oxygen exposure studies WDOE (2002) concluded salmonid growth rates decreased less than 10% at dissolved oxygen concentrations of 8 mg/L or more, less than 20% at 7 mg/L, and generally less than 22% at 5 - 6 mg/L. Herrmann (1958) found that the mean percentage of weight gain in juvenile coho held at constant dissolved oxygen concentrations was 7.2% around 2 mg/L, 33.6% at 3 mg/L, 55.8% near 4 mg/L, and 67.9% at or near 5 mg/L. In a laboratory study Fischer (1963) found that the growth rates of juvenile coho exposed to constant oxygen concentrations ranging from 2.5 to 35.5 mg/L (fed to satiation, temperature at approximately 18 °C) dramatically

decreased with decreases in the oxygen concentration below 9.5 mg/L (air saturation level). WDOE (2002) concludes that a monthly or weekly average concentration of 9 mg/L, and a monthly average of the daily minimum concentrations should be at or above 8 - 8.5 mg/L to have a negligible effect (5% or less) on growth and support healthy growth rates.

Food conversion efficiency is related to dissolved oxygen levels and the process becomes less efficient when oxygen concentrations are below 4 - 4.5 mg/L (ODEQ 1995b). Bjornn and Reiser (1991) state that growth, food conversion efficiency, and swimming performance are adversely affected when dissolved oxygen concentrations are <5 mg/L. The USEPA (1986a) reviewed growth data from a study conducted by Warren et al. (1973) where tests were conducted at various temperatures to determine the growth of coho and Chinook. USEPA cites that, with the exception of tests conducted at 22 °C, the results supported the idea that the effects of low dissolved oxygen become more severe at higher temperatures.

Brett and Blackburn (1981) performed a laboratory study to determine the growth rate and food conversion efficiency of young coho and sockeye salmon fed full rations. Tests were performed at dissolved oxygen concentrations ranging from 2 to 15 mg/L at a constant temperature of 15°C, the approximate optimum temperature for growth of Pacific Salmon. Both species showed a strong dependence of growth on the environmental oxygen concentrations when levels were below 5 mg/L. For coho, zero growth was observed at dissolved oxygen concentrations of 2.3 mg/L. The mean value for maximum coho growth occurred at 4 mg/L, and at dissolved oxygen concentrations above this level growth did not appear to be dependant on the dissolved oxygen. Sockeye displayed zero growth at oxygen levels of 2.6 mg/L, and reached the zone of independence (growth not dependant on dissolved oxygen levels) at 4.2 mg/L. Brett and Blackburn (1981) conclude that the critical inflection from oxygen dependence to independence occurs at 4 - 4.2 mg/L for coho and sockeye.

Herrmann et al. (1962) studied the influence of various oxygen concentrations on the growth of age 0 coho salmon held at 20 °C. Coho were held in containers at a constant mean dissolved oxygen level ranging from 2.1 - 9.9 mg/L and were fed full rations. The authors concluded that oxygen concentrations below 5 mg/L resulted in a sharp decrease in growth and food consumption. A reduction in the mean oxygen levels from 8.3 mg/L to 6 and 5 mg/L resulted in slight decreases in food consumption and growth. Weight gain in grams per gram of food consumed was slightly depressed at dissolved oxygen concentrations near 4 mg/L, and were markedly reduced at lower concentrations. At oxygen levels of 2.1 and 2.3 mg/L, many fish died and the surviving fish lost weight and consumed very little food.

USEPA (1986a) calculated the median percent reduction in growth rate of Chinook and coho salmon fed full rations at various dissolved oxygen concentrations. They calculated no reduction in growth at dissolved oxygen concentrations of 8 and 9 mg/L, and a 1% reduction in growth at 7 mg/L for both species. At 6 mg/L Chinook and coho growth were reduced by 7% and 4% respectively. Dissolved oxygen levels of 4 mg/L result in a

29% reduction in growth for Chinook salmon and 21% reduction in growth for coho. At 3 mg/L there was a 47% decrease in Chinook growth and a 37% reduction in coho growth. USEPA (1986a) states that due to the variability inherent in growth studies the reductions in growth rates seen above 6 mg/L are not usually statistically significant, while reductions in growth at dissolved oxygen levels below 4 mg/L are considered severe.

2.2.7.3 Avoidance and Preference

Salmonids have been reported to actively avoid areas with low dissolved oxygen concentrations, which is likely a useful protective mechanism that enhances survival (Davis 1975). Field and laboratory studies have found that avoidance reactions in juvenile salmonids consistently occur at concentrations of 5 mg/L and lower, and there is some indication that avoidance is triggered at concentrations as high as 6 mg/L. Therefore these dissolved oxygen levels should be considered a potential barrier to the movement and habitat selection of salmonids (WDOE 2002).

Spoor (1990) performed a laboratory study on the distribution of fingerling brook trout in dissolved oxygen concentration gradients. Sixteen gradients between 1 and 8.9 mg/L were used for the study to determine what level of dissolved oxygen is preferred by the brook trout. It was found that in the absence of a gradient with dissolved oxygen concentrations at 6 mg/L or more throughout the system, the fish moved freely without showing preference or avoidance. Movement from low to higher oxygen concentrations were noted throughout the study. Fish moved away from water with dissolved oxygen concentrations from 1 - 1.9 mg/L within one hour, moved away from water with dissolved oxygen concentrations of 2 - 2.9 mg/L within 1 - 2 hours, and moved away more slowly from concentrations of 3 - 3.9 mg/L. From his study, Spoor (1990) concluded that brook trout will avoid oxygen concentrations below 4 mg/L, and preferred oxygen levels of 5 mg/L or higher.

Whitmore et al. (1960) performed studies with juvenile coho and Chinook salmon to determine their avoidance reaction to dissolved oxygen concentration of 1.5, 3, 4.5, and 6 mg/L at variable river water temperatures. Juvenile Chinook salmon showed marked avoidance of oxygen concentrations near 1.5, 3, and 4.5 mg/L in the summer at mean temperatures ranging from 20.7 - 22.8°C, but no avoidance to levels near 6 mg/L at a mean temperature of 18.4°C. Chinook did not show as strong an avoidance to these oxygen levels in the fall when water temperatures were lower, ranging from 11.8 -13.2°C. Chinook showed little avoidance of dissolved oxygen concentrations near 4.5 mg/L during the fall, and no avoidance to concentrations near 6 mg/L. In all cases avoidance became progressively larger with reductions in the oxygen concentration below 6 mg/L. Seasonal differences of avoidance are most likely due to differences in water temperature. At temperatures ranging from 18.4 - 19°C juvenile coho salmon showed some avoidance to all of the above oxygen concentrations, including 6 mg/L. Their behavior was more erratic than that of Chinook, and their avoidance of concentrations near 4.5 mg/L and lower was not as pronounced at corresponding temperatures. The juvenile coho often started upon entering water with low dissolved

oxygen and then darted around until they found their way out of the experimental channel.

USEPA (1986a) performed a literature review and cites the effects of various dissolved oxygen concentrations on salmonid life stages other than embryonic and larval (Table 16). These effects range from no impairment at 8 mg/L to acute mortality at dissolved oxygen levels below 3 mg/L.

Table 16: Dissolved oxygen concentrations and their effects on salmonid life stages other than embryonic and larval

Level of Effect	Water Column DO (mg/L)
No Production Impairment	8
Slight Production Impairment	6
Moderate Production Impairment	5
Severe Production Impairment	4
Limit to Avoid Acute Mortality	3

Source: USEPA 1986a

2.2.8 Lethality

Salmonid mortality begins to occur when dissolved oxygen concentrations are below 3 mg/L for periods longer than 3.5 days (USEPA 1986a). A summary of various field study results by WDOE (2002) reports that significant mortality occurs in natural waters when dissolved oxygen concentrations fluctuate the range of 2.5 - 3 mg/L. Long-term (20 - 30 days) constant exposure to mean dissolved oxygen concentrations below 3 - 3.3 mg/L is likely to result in 50% mortality of juvenile salmonids (WDOE 2002). According to a short-term (1 - 4 hours) exposure study by Burdick et al. (1954, as cited by WDOE, 2002), in warm water (20 - 21°C) salmonids may require daily minimum oxygen levels to remain above 2.6 mg/L to avoid significant (50%) mortality. From these and other types of studies, WDOE (2002) concluded that juvenile salmonid mortality can be avoided if daily minimum dissolved oxygen concentration remain above 3.9 mg/L, and the monthly or weekly average of minimum concentrations remains above 4.6 mg/L.

2.3 Effects of High Total Dissolved Gas Concentrations on Salmonids

High levels of total dissolved gas (TDG), including dissolved oxygen, can be harmful to salmonids and other fish and result in "gas bubble disease". This occurs when dissolved gases in their circulatory system come out of solution and form bubbles which block the flow of blood through the capillary vessels (USEPA 1986b). There are several ways TDG supersaturation can occur, including excessive algal photosynthesis which can create supersaturated dissolved oxygen conditions (USEPA 1986b). Thus, to protect salmonids and other freshwater fish the USEPA has set criteria for TDG stating that levels should not exceed 110% of the saturation value.

Numerous studies have been conducted to determine the mortality rate of salmonids exposed to various levels of TDG. Mesa et al. (2000) conducted laboratory experiments on juvenile Chinook and steelhead, exposing them to different levels of TDG and found no fish died when held at 110% TDG for up to 22 days. When fish were exposed to 120% TDG, 20% of juvenile Chinook died within 40 to 120 hours while 20% of juvenile

steelhead died within 20 to 35 hours. At TDG levels of 130% Chinook mortality reached 20% after 3 to 6 hours and steelhead mortality was 20% after 5 to 7 hours. Gale et al. (2001) held adult female spring Chinook at mean TDG levels ranging from 114.1% to 125.5% and found the time to first mortality ranged from 10 to 68 hours.

USEPA (1986b) discusses various studies on the effects of TDG on salmonids. The following studies are all cited from the USEPA 1986 water quality criteria document. Bouck et al. (1975) found TDG levels of 115% and above to be acutely lethal to most species of salmonids, and levels of 120% TDG are rapidly lethal to all salmonids. Conclusions drawn from Ebel et al. (1975) and Rulfison and Abel (1971) include the following:

- Adult and juvenile salmonids confined to shallow water (1 m) with TDG levels above 115% experience substantial levels of mortality.
- Juvenile salmonids exposed sublethal levels TDG supersaturation are able to recover when returned to normally saturated water, while adults do not recover and generally die.

CHAPTER 3. AMMONIA

3.1 Introduction

According to the USEPA (1986b, p.17), acute concentrations of ammonia can cause loss of equilibrium, hyperexcitability, increased breathing, cardiac output and oxygen uptake, and, in extreme cases, convulsions, coma, and death of fish. Lower concentrations of ammonia can result in reduced hatching success, reduced growth and morphological development, and pathologic changes in tissues of gills, livers, and kidneys.

The information in the following sections was extracted from the USEPA document titled: 1999 Update of Ambient Water Quality Criteria for Ammonia. The information presented applies to salmonids in general.

3.2 Ammonia Speciation

Ammonia in water exists primarily in two forms, un-ionized ammonia (NH₃) and ammonium ion (NH₄⁺) (USEPA 1999b, p.2). The fraction of each of these two forms, or ammonia speciation, varies markedly with temperature and pH (USEPA 1999b, p.2). The pH-dependence of the relative amounts of un-ionized ammonia and ammonium ion at 25°C are presented in Figure 1 below (USEPA 1999b, p.2). Ammonia speciation also depends on ionic strength, although in freshwater this effect is much smaller than the effects of temperature and pH (USEPA 1999b, p.3)




Figure 1: Chemical Speciation of Ammonia

Source: USEPA 1999b, p.3

3.3 Ammonia Toxicity

These speciation relationships are important to ammonia toxicity because un-ionized ammonia is much more toxic than ammonium ion. The importance of un-ionized ammonia was first recognized when it was observed that increased pH caused total ammonia to appear to be much more toxic (Chipman 1934; Wuhrmann and Woker 1948). It is not surprising that un-ionized ammonia is the more toxic form, because it is a neutral molecule and thus is able to diffuse across the epithelial membranes of aquatic organisms much more readily than the charged ammonium ion. Ammonia is unique among regulated pollutants because it is an endogenously produced toxicant that organisms have developed various strategies to excrete, which is in large part by passive diffusion of un-ionized ammonia from the gills. High external un-ionized ammonia concentrations reduce or reverse diffusive gradients and cause the buildup of ammonia in gill tissue and blood (USEPA 1999b, p.3).

Because of the importance of un-ionized ammonia, it became a convention in the scientific literature to express ammonia toxicity in terms of un-ionized ammonia, and water quality criteria and standards followed this convention. However, there are reasons to believe that ammonium ion can contribute significantly to ammonia toxicity under some conditions. Observations that ammonia toxicity is relatively constant when expressed in terms of un-ionized ammonia come mainly from toxicity tests conducted at pH>7.5. At lower pH, toxicity varies considerably when expressed in terms of unionized ammonia and under some conditions is relatively constant in terms of ammonium ion (Erickson 1985). Also, studies have established that mechanisms exist for the transport of ammonium ion across gill epithelia (Wood 1993), so this ion might contribute significantly to ammonia exchange at gills and affect the buildup of ammonia in tissues if its external concentration is sufficiently high. Thus, the very same arguments employed for the importance of un-ionized ammonia can also be applied in some degree to ammonium ion. This is not to say that ammonium ion is as toxic as unionized ammonia, but rather that, regardless of its lower toxicity, it can still be important because it is generally present in much greater concentrations than un-ionized ammonia (USEPA 1999b, p.3,4).

3.4 Ammonia Criteria

The USEPA has utilized the above information to create pH-dependant acute and pH-and temperature-dependent chronic criterion for total ammonia (NH₃ and NH₄⁺) as nitrogen in freshwater (Tables 17, 18, 19).

Table 17: pH-Dependent Values of the Criterion Maximum Concentration (CMC) of Total Ammonia as Nitrogen (mg N/L) in Freshwater when Salmonids are Present

	Acute Criterion ¹							
pН	CMC Total NH3 mgN/L	pН	CMC Total NH3 mgN/L					
6.5	32.6	7.8	8.11					
6.6	31.3	7.9	6.77					
6.7	29.8	8.0	5.62					
6.8	28.1	8.1	4.64					
6.9	26.2	8.2	3.83					
7.0	24.1	8.3	3.15					
7.1	22.0	8.4	2.59					
7.2	19.7	8.5	2.14					
7.3	17.5	8.6	1.77					
7.4	15.4	8.7	1.47					
7.5	13.3	8.8	1.23					
7.6	11.4	8.9	1.04					
7.7	9.65	9.0	0.885					

¹ The one-hour average concentration of total ammonia nitrogen (NH₃ and NH₄⁺) should not exceed this value more than once every 3 years.

Source: USEPA 1999b, p.86

Table 18: Temperature and pH-Dependent Values of the Criterion Continuous Continuation (CCC) for Total Ammonia as Nitrogen (mg N/L) in Freshwater when Fish Early Life Stages are Present

	Chronic Criterion ¹										
	CCC for Fish Early Life Stages Present, mg N/L										
		Temperature, C									
1	рH	0	14	16	18	20	22	24	26	28	30
6	3.5	6.67	6.67	6.06	5.33	4.68	4.12	3.62	3.18	2.80	2.46
6	6.6	6.57	6.57	5.97	5.25	4.61	4.05	3.56	3.13	2.75	2.42
6	6.7	6.44	6.44	5.86	5.15	4.52	3.98	3.50	3.07	2.70	2.37
6	8.6	6.29	6.29	5.72	5.03	4.42	3.89	3.42	3.00	2.64	2.32
6	6.9	6.12	6.12	5.56	4.89	4.30	3.78	3.32	2.92	2.57	2.25
7	7.0	5.91	5.91	5.37	4.72	4.15	3.65	3.21	2.82	2.48	2.18
7	7.1	5.67	5.67	5.15	4.53	3.98	3.50	3.08	2.70	2.38	2.09
7	7.2	5.39	5.39	4.90	4.31	3.78	3.33	2.92	2.57	2.26	1.99
7	7.3	5.08	5.08	4.61	4.06	3.57	3.13	2.76	2.42	2.13	1.87
7	7.4	4.73	4.73	4.30	3.78	3.32	2.92	2.57	2.26	1.98	1.74
7	7.5	4.36	4.36	3.97	3.49	3.06	2.69	2.37	2.08	1.83	1.61
7	7.6	3.98	3.98	3.61	3.18	2.79	2.45	2.16	1.90	1.67	1.47
7	7.7	3.58	3.58	3.25	2.86	2.51	2.21	1.94	1.71	1.50	1.32
7	7.8	3.18	3.18	2.89	2.54	2.23	1.96	1.73	1.52	1.33	1.17
7	7.9	2.80	2.80	2.54	2.24	1.96	1.73	1.52	1.33	1.17	1.03
8	3.0	2.43	2.43	2.21	1.94	1.71	1.50	1.32	1.16	1.02	0.897
8	3.1	2.10	2.10	1.91	1.68	1.47	1.29	1.14	1.00	0.879	0.773
8	3.2	1.79	1.79	1.63	1.43	1.26	1.11	0.973	0.855	0.752	0.661
8	3.3	1.52	1.52	1.39	1.22	1.07	0.941	0.827	0.727	0.639	0.562
8	3.4	1.29	1.29	1.17	1.03	0.906	0.796	0.700	0.615	0.541	0.475
8	3.5	1.09	1.09	0.990	0.870	0.765	0.672	0.591	0.520	0.457	0.401
8	3.6	0.920	0.920	0.836	0.735	0.646	0.568	0.499	0.439	0.386	0.339
8	3.7	0.778	0.778	0.707	0.622	0.547	0.480	0.422	0.371	0.326	0.287
8	3.8	0.661	0.661	0.601	0.528	0.464	0.408	0.359	0.315	0.277	0.244
8	3.9	0.565	0.565	0.513	0.451	0.397	0.349	0.306	0.269	0.237	0.208
6	9.0	0.486	0.486	0.442	0.389	0.342	0.300	0.264	0.232	0.204	0.179

¹ The thirty-day average concentration of total ammonia (NH₃ and NH₄⁺) should not exceed this value more than once every three years. Additionally, the highest four day average within the thirty-day period should not exceed 2.5 times the CCC (USEPA 1999b, p.87).

Table 19: Temperature and pH-Dependent Values of the Criterion Continuous Continuation (CCC) for Total Ammonia as Nitrogen (mg N/L) in Freshwater when Fish Early Life Stages are Absent

Chronic Criterion ¹										
CCC for Fish Early Life Stages Absent, mg N/L										
	Temperature									
pН	0-7	8	9	10	11	12	13	14	15*	16*
6.5	10.8	10.1	9.51	8.92	8.36	7.84	7.35	6.89	6.46	6.06
6.6	10.7	9.99	9.37	8.79	8.24	7.72	7.24	6.79	6.36	5.97
6.7	10.5	9.81	9.20	8.62	8.08	7.58	7.11	6.66	6.25	5.86
6.8	10.2	9.58	8.98	8.42	7.90	7.40	6.94	6.51	6.10	5.72
6.9	9.93	9.31	8.73	8.19	7.68	7.20	6.75	6.33	5.93	5.56
7.0	9.60	9.00	8.43	7.91	7.41	6.95	6.52	6.11	5.73	5.37
7.1	9.20	8.63	8.09	7.58	7.11	6.67	6.25	5.86	5.49	5.15
7.2	8.75	8.20	7.69	7.21	6.76	6.34	5.94	5.57	5.22	4.90
7.3	8.24	7.73	7.25	6.79	6.37	5.97	5.60	5.25	4.92	4.61
7.4	7.69	7.21	6.76	6.33	5.94	5.57	5.22	4.89	4.59	4.30
7.5	7.09	6.64	6.23	5.84	5.48	5.13	4.81	4.51	4.23	3.97
7.6	6.46	6.05	5.67	5.32	4.99	4.68	4.38	4.11	3.85	3.61
7.7	5.81	5.45	5.11	4.79	4.49	4.21	3.95	3.70	3.47	3.25
7.8	5.17	4.84	4.54	4.26	3.99	3.74	3.51	3.29	3.09	2.89
7.9	4.54	4.26	3.99	3.74	3.51	3.29	3.09	2.89	2.71	2.54
8.0	3.95	3.70	3.47	3.26	3.05	2.86	2.68	2.52	2.36	2.21
8.1	3.41	3.19	2.99	2.81	2.63	2.47	2.31	2.17	2.03	1.91
8.2	2.91	2.73	2.56	2.40	2.25	2.11	1.98	1.85	1.74	1.63
8.3	2.47	2.32	2.18	2.04	1.91	1.79	1.68	1.58	1.48	1.39
8.4	2.09	1.96	1.84	1.73	1.62	1.52	1.42	1.33	1.25	1.17
8.5	1.77	1.66	1.55	1.46	1.37	1.28	1.20	1.13	1.06	0.990
8.6	1.49	1.40	1.31	1.23	1.15	1.08	1.01	0.951	0.892	0.836
8.7	1.26	1.18	1.11	1.04	0.976	0.915	0.858	0.805	0.754	0.707
8.8	1.07	1.01	0.944	0.885	0.829	0.778	0.729	0.684	0.641	0.601
8.9	0.917	0.860	0.806	0.756	0.709	0.664	0.623	0.584	0.548	0.513
9.0	0.790	0.740	0.694	0.651	0.610	0.572	0.536	0.503	0.471	0.442

¹ The thirty-day average concentration of total ammonia (NH₃ and NH₄⁺) should not exceed this value more than once every three years. Additionally, the highest four day average within the thirty-day period should not exceed 2.5 times the CCC.

Source: USEPA 1999b, p.88

CHAPTER 4. pH

4.1 Introduction

The pH of freshwater streams is important for adult and juvenile salmonid development. Chronic effects from low pH can occur at levels that are not toxic to adult fish but that impair reproduction including altered spawning behavior, reduced egg viability, decreased hatchability, and reduced survival of the early life stages (Jordahl and Benson 1987). The early life stages of salmonid development are most vulnerable to low pH (Jordahl and Benson 1987). Chronic high pH levels in freshwater streams can decrease activity levels of salmonids, create stress responses, decrease or cease feeding, and lead to a loss of equilibrium (Murray and Ziebell 1984; Wagner et al. 1997). Additionally, high temperatures can exacerbate the effects of high pH levels on salmonids (Wagner et al. 1997). If pH reaches extremely low or high levels, death can occur (Wagner et al. 1997).

Literature reviewed for this analysis included numerous laboratory and field studies. The following information applies to salmonids in general.

4.2 Effects of High pH

Wagner et al. (1997) conducted laboratory and field studies and found that pH values of greater than 9.4 will result in the death of rainbow trout, especially at temperatures ranging from 19-22 C. Fresh water pH values of 9.0 or greater resulted in significant stress responses in rainbow trout.

Wilkie and Wood (1996) found that Lahontan cutthroat trout exposed to high pH waters (9.4) permanently lowered their rate of nitrogenous waste production to avoid the potentially toxic build-up of internal ammonia. However, rainbow trout, kokanee, and brown trout were unable to adapt to the high pH and died.

Murray and Ziebell (1984) found that rainbow trout are not able to acclimate to pH levels of 10.0 or higher and that their ability to tolerate pH above 9.0 depends on the rate of acclimation. Gradual acclimation (0.2 to 0.4 of a pH unit/day) allowed rainbow trout to acclimate to a pH of 9.8 and continue feeding, although they showed signs of distress and their activity was greatly reduced by the end of 4 days when the pH reached 9.9 (Table 20). The maximum pH tolerated before fish began dying was 10.2.

Rapid acclimations tests conducted by Murray and Ziebell (1984) yeilded the following results:

Rainbow trout mortalities were 40% or greater in preliminary acclimation tests in which pH was increased to 9.6 and 9.7 in 3 and 5 hours. These results were comparable to previous shock tests (unpublished data). Consequently, in later experiments, acclimation time was increased to 6 hours and pH values were lowered to 9.3 and 9.5.

Trout in the pH 9.5 experiments exhibited adverse reactions and mortalities were similar to those seen in preliminary tests at pH 9.6 and 9.7. All fish began to show marked signs of stress within 12 hours, and within 24 hours the mortalities in replicated experiments were 30, 40, and 50% respectively. At 49 hours the last deaths occurred that brought mortalities to 50% in each replicate. All remaining living fish were distressed and did not feed. After 72 hours had elapsed, the survivors resumed feeding and their condition improved until the experiments were terminated at 120 hours.

In the pH 9.3, 6-hour acclimation experiments trout exhibited only minor adverse reactions. The primary behavioral changes were a decrease in swimming activity and a temporary loss of appetite. After 48 hours all fish resumed normal feeding and became progressively more active. No mortalities occurred in any of the replicated experiments, and all fish behaved normally when the experiments were terminated at 120 hours Murray and Ziebell (1984).

Table 20: Reactions of 10 rainbow trout to various pH levels during gradual acclimation

experiments (0.2 to 0.4 of a pH unit/day)

Day	pH Range	Reactions and Condition of Trout
1	8.6-8.9	Normal
2	8.9-9.2	Activity decreased but feeding normal
3	9.2-9.7	Activity further decreased but feeding continued
4	9.7-9.9	Minor distress shown but feeding continued
5	9.9-10.3	Some fish lost equilibrium at 10.0, and feeding ceased. Loss of equilibrium increased at 10.1 and eyes of some fish developed corneal opacities; 50% of fish lost equilibrium at pH 10.2 and mortality was 60% at pH 10.3

Source: Murray and Ziebell (1984)

4.3 Effects of Low pH

"Chronic effects of low pH on fish populations may occur at pH levels that are not toxic to adult fish but that impair reproduction, and ultimately lead to population extinction (Jordahl and Benson 1987)." A study was conducted by Weiner et al. (1986) to determine the effects of low pH on the reproductive success of rainbow trout. It was determined that exposure of adult salmonids to pH values below 5.5 negatively effected reproduction. Adult rainbow trout were exposed to pH 4.5, 5.0, 5.5, and 6.5-7.1 during the final 6 weeks of reproductive maturation. Weiner et al. found that pH values of 5.5 and below impaired the creation of eggs in females and sperm in males.

Jordahl and Benson (1987) report that reproductive failure occurred in adult brook trout due to low pH in a freshwater stream with pH levels ranging from 5.0-5.8, while trout in a reference stream with pH ranging from 6.1-7.2 did not experience reproductive failure. Additionally, brook trout were absent from a highly acidic freshwater stream with pH ranging from 4.7-5.4 leading Jordahl and Benson to conclude that breeding females may avoid acidic tributaries.

In addition to effecting adult salmonids, highly acidic freshwater (low pH) can have a detrimental effect on eggs and juvenile salmonids. Weiner et al. (1986) determined that juvenile rainbow trout mortality was greatly increased at pH levels of 5.5 and below, and that no eggs survived when exposed to pH levels below 4.5. Hulsman and Powels (1983) found the mortality of rainbow trout yolk-sac larvae approached 100% within 5 days of exposure to pH 4.6 and 5.4, whereas exposure to pH 6.0 resulted in less than 3% mortality.

Jordahl and Benson (1987) conducted a study to determine the effect of low pH on juvenile brook trout survival and found that survival rates were highest in a freshwater stream with pH values ranging from 6.1-7.2 and lower in acidified streams with pH levels of 4.7-5.8. At pH values of 5.0 and lower, growth was retarded and the development of yolk-sac larvae was considerably prolonged. Additionally, larval activity was depressed, pigmentation was reduced, and incomplete hatching was observed in streams with low pH values of 4.7-5.8, but not in the stream with pH ranging from 6.1-7.2. Jordahl and Benson concluded that mean pH values of 5.0-5.4 can cause acid stress on developing juveniles, while pH levels from 6.1-7.2 are above ranges that negatively effect juvenile brook trout development and survival.

REFERENCES

- Armour, C.L. 1991. Guidance for evaluating and recommending temperature regimes to protect fish. U.S. Fish and Wildlife Service Biological Report 90(22). 13 pp.
- Bell, M.C.. 1986. Fisheries handbook of engineering requirements and biological criteria. Fish Passage Development and Evaluation Program. U.S. Army Corps of Engineers. 209pp.
- Bjornn, T. and D. Reiser. 1991. Habitat requirements of salmonids in streams. In Meehan, W. ed., <u>Influences of Forest and Rangeland Management on Salmonids Fishes and Their Habitat</u>. American Fisheries Society Special Publication 19. pp. 83-138.
- Brett, J.R. 1952. Temperature Tolerance in Young Pacific Salmon, Genus Oncorhynchus. Pacific Biological Station, and Department of Zoology, University of Toronto. J. Fish Res. Board Can. 9(6): 265-308 + appendices.
- Brett, J. R., and J.M. Blackburn. 1981. Oxygen requirements for growth of young coho (*Oncorhynchus kisutch*) and sockeye (*O. nerka*) salmon at 15C. Can. J. Fish. Aquat. Sci. 38:399-404.
- Brungs, W.A., and B.R. Jones. 1977. Temperature criteria for freshwater fish: Protocol and procedures. Environmental Research Laboratory, Duluth, MN. U.S. Environmental Protection Agency. EPA-600/3-77-061. 136pp.
- California Department of Fish and Game (CDFG). 2001. California's Living Marine Resources: A Status Report. Leet, W.S., C.M. Dewees, R. Klingbeil, and E.J. Larson [eds.]. The Resources Agency. Sacramento, CA. December 2001. 552pp. + appendices. Available at: http://www.dfg.ca.gov/mrd/status/steelhead-rainbow_trout.pdf. Website accessed on July 21, 2004.
- California Department of Water Resources (CDWR). 1988. Water temperature effects on Chinook salmon (*Oncorhynchus tshawytscha*) with emphasis on the Sacramento River: a literature review. Northern District Office Report, Red Bluff, California. 42pp.
- Cobel, D.W. 1961. Influence of water exchange and dissolved oxygen in redds on survival of steelhead trout embryos. Transactions of the American Fisheries Society. 90:469-474.
- Dahlberg, M.L., D.L. Shumway, and P. Doudoroff. 1968. Influence of dissolved oxygen and carbon dioxide on swimming performance of largemouth bass and coho salmon. Journal of the Fisheries Research Board of Canada. 25(1):49-70.

- Davis, G.E., J. Foster, C.E. Warren, and P. Doudoroff. 1963. The influence of oxygen concentration on the swimming performance of juvenile pacific salmon at various temperatures. Transactions of the American Fisheries Society. 92:111-124.
- Davis, J.C. 1975. Minimal dissolved oxygen requirements of aquatic life with emphasis on canadian species: a review. Journal of the Fisheries Research Board of Canada. 32:2295-2332.
- Dickerson, H.W. and D.L. Dower. 1995. *Ichthyophthirius multifiliis* and *Cryptocaryon irritans* (Phylum Ciliophora). In Woo, P.T.K. ed., <u>Fish Disease and Disorders</u>. ISBN 085198 823 7; p.181-227.
- Durborow, R.M., A.J. Mitchell, and M.D. Crosby. 1998. Ich (White Spot Disease). Southern Regional Aquaculture Center, Delta Research and Extension Center, Mississippi State University. Stoneville, Mississippi. 6pp.
- Eddy, R.M. 1971. The influence of dissolved oxygen concentration and temperature on the survival and growth of Chinook salmon embryos and fry. Masters of Science Thesis. Oregon State University. Corvallis, Oregon. 45pp.
- Elliott, J. 1981. Some aspects of thermal stress on freshwater teleosts. In Pickering, A.D. ed., <u>Stress and Fish</u>: Academic Press, London, p. 209-245.
- Fischer, R.J. 1963. Influence of Oxygen Concentration and of its Diurnal Fluctuations on the Growth of Juvenile Coho Salmon. Masters of Science Thesis. Oregon State University. Corvallis, Oregon. 48pp.
- Gale, W.L., A.G. Maule, A. Postera, and M. Peter-Swihart. 2001. Effects of Supersaturated Water on Reproductive Success of Adult Salmonids. Bonneville Power Administration project no. 2000-58. December 2, 2001. 33 pp.
- Garside, E.T. 1966. Effects of oxygen in relation to temperature on the development of embryos of brook trout and rainbow trout. J. Fish. Res. Bd. Canada 23(8)1121-1134.
- Hallock, R.J., R.F. Elwell, and D.H. Fry, Jr. 1970. Migrations of adult king salmon *Oncorhynchus tshawytsca* in the San Joaquin Delta as demonstrated by the use of sonic tags. California Department of Fish and Game, Fish Bulletin 151. 92pp.
- Hassler, T.J. 1987. Species Profiles: Life Histories and Environmental Requirements of Coastal Fishes and Invertebrates (Pacific Southwest). Coho Salmon. U.S. Fish and Wildlife Service Biological Report 82(11.70). U.S. Army Corps of Engineers, TR EL-82-4. 19pp.

- Herrmann, R.B. 1958. Growth of juvenile coho salmon at various concentrations of dissolved oxygen. Masters of Science Thesis. Oregon State University. Corvallis, Oregon. 82pp.
- Herrmann, R.B., C.E. Warren, and P. Doudoroff. 1962. Influence of oxygen concentration on the growth of juvenile coho salmon. Transactions of the American Fisheries Society. 91:155-167.
- Holt, R.A., J.E. Sanders, J.L. Zinn, J.L. Fryer, and K.S. Pilcher. 1975. Relation of Water Temperature to *Flexibacter columnaris* Infection in Steelhead Trout (*Salmo gairdneri*), Coho (*Oncorhynchus kisutch*) and Chinook (*O. tshawytscha*) Salmon. J. Fish. Res. Board Can. 32:1553-1559.
- Hulsman P.F. and P.M. Powles. 1983. Mortality of Walleye Eggs and Rainbow Trout Yolk-Sac Larvae in Low-pH Waters of the LaCloche Mountain Area, Onterio. Transactions of the American Fisheries Society. 112:680-688.
- Jordahl and Benson, 1987. Effect of Low pH on Survival of Brook Trout Embroys and Yolk-Sac Larvae in West Virginia Streams. Transactions of the American Fisheries Society. 116:807-816.
- Konecki, T.C., C.A. Woody, and T.P. Quinn. 1995a. Temperature preference in two populations of juvenile coho salmon, *Oncorhynchus kisutch*. Northwest Science. 69 (2): 417-421.
- Konecki, T.C., C.A. Woody, and T.P. Quinn. 1995b. Critical thermal maxima of coho salmon (*Oncorhynchus kisutch*) fry under field and laboratory acclimation regimes. Can. J. Zool. 73:993-996.
- Leitritz, E. and R.C. Lewis. 1976. Trout and salmon culture. California Department of Fish and Game. Fish Bulletin 164. 197pp.
- Ligon, F., A. Rich, G. Rynearson, D. Thornburgh, and W. Trush. 1999. Report of the Scientific Review Panel on California Forest Practice Rules and Salmonid Habitat: Prepared for the Resource Agency of California and the National Marine Fisheries Sacramento, Calif. 92pp. + appendices.
- Marine, K.R., and J.J. Cech. 2004. Effects of High Water Temperature on Growth, Smoltification, and Predator Avoidance in Juvenile Sacramento River Chinook Salmon. North American Journal of Fisheries Management 24:198-210.
- McMahon, T.E. 1983. Habitat suitability index models: Coho salmon. U.S. Department of Interior, Fish and Wildlife Service. FWS/OBS-82/10.49. 29pp.

- Mesa, M.G., L.K. Weiland, and A.G. Maule. 2000. Progression and Severity of Gas Bubble Trauma in Juvenile Salmonids. Transactions of the American Fisheries Society. 129:174-185.
- Murray and Ziebell. 1984. Acclimation of Rainbow Trout to High pH to Prevent Stocking Mortality in Summer. The Progressive Fish-Culturist. 46(3):176-179.
- Murray, C.B., and J.D. McPhail. 1988. Effect of incubation temperature on the development of five species of Pacific salmon (*Oncorhynchus*) embryos and alevins. Can. J. Zool. 66:266-273.
- Myrick C.A., and J.J. Cech. 2001. Temperature Effects on Chinook Salmon and Steelhead: a Review Focusing on California's Central Valley Populations. Bay-Delta Modeling Forum. Technical Publication 01-1. 57pp.
- Nielsen, J.L., T.E. Lisle, and V. Ozaki. 1994. Thermally stratified pools and their use by steelhead in Northern California streams. Transactions of the American Fisheries Society, 123:613-626. Available at: http://www.humboldt.edu/~storage/ pdfmill/Batch_9/thermally.pdf >. Website accessed on August 27, 2004.
- Oregon Department of Environmental Quality (ODEQ). 1995a. Temperature: 1992-1994 Water quality standards review. Final Issue Paper. 122pp. Available at: http://www.fishlib.org/Bibliographies/waterquality.html>. Website accessed on August 20, 2004.
- Oregon Department of Environmental Quality (ODEQ). 1995b. Dissolved Oxygen: 1992-1994 Water quality standards review. Final Issue Paper. 166pp. Available at: http://www.fishlib.org/Bibliographies/waterquality.html>. Website accessed August 20, 2004.
- Pacha, R.E. and E. J. Ordal. 1970. Myxobacterial diseases of salmonids. In Snieszko, S.F. ed., <u>A symposium on diseases of fishes and shellfishes</u>. Special Publication No. 5, American Fisheries Society, Washington, D.C. p.243-257.
- Piper, R.G., I.B. McElwain, L.E. Orme, J.P. McCraren, L.G. Fowler, and J.R. Leonard. 1982. Fish hatchery management. U.S. Department of Interior, Fish and Wildlife Service, Washington D.C. 517 pp.
- Reutter, J.M. and C.E. Herdendorf. 1974. Laboratory Estimates of the Seasonal Final Temperature Preferenda of Some Lake Erie Fish. Presented at the 17th Conference for the International Association for Great Lakes Research. Center for Lake Erie Area Research, Ohio State University. Columbus, Ohio. Clear Technical Report No. 20. 13pp. + figures & tables.

- Shumway, D.L., C.E. Warren, and P. Doudoroff. 1964. Influence of oxygen concentration and water movement on the growth of steelhead trout and coho salmon embryos. Transactions of the American Fisheries Society. 93:342-356.
- Silver, S.J., C.E. Warren, P. Doudoroff. 1963. Dissolved oxygen requirements of developing steelhead trout and Chinook salmon embryos at different water velocities. Transactions of the American Fisheries Society. 92(4):327-343.
- Spence, B.C., G.A. Lomicky, R.M. Hughes, and R.P. Novitzki. 1996. An ecosystem approach to salmonid conservation. TR-4501-96-6057. ManTech Environmental Research Services Corp., Corvallis, OR (Available from the National Marine Fisheries Service, Portland, OR). 356pp.
- Spoor, W.A. 1990. Distribution of fingerling brook trout, *Salvelinus fontinalis* (Mitchill), in dissolved oxygen concentration gradients. J. Fish. Biol. 36:363-373.
- Strange, J. 2006. Adult Chinook Salmon Migration in the Klamath River Basin: 2005 Sonic Telemetry Study Report (DRAFT). Yurok Tribal Fisheries Program, and School of Aquatic and Fishery Sciences- University of Washington in collaboration with Hoopa Valley Tribal fisheries. April 2006. 93pp.
- Sullivan K., D.J. Martin, R.D. Cardwell, J.E. Toll, and S. Duke. 2000. An analysis of the effects of temperature on salmonids of the Pacific Northwest with implications for selecting temperature criteria. Sustainable Ecosystems Institute. Portland, OR. 147 pp. Available at: http://www.sei.org/pub.html#reports>. Website accessed on June 11, 2004.
- Thomas, R.E., J.A. Gharrett, M.G. Carls, S.D. Rice, A. Moles, and S. Korn. 1986. Effects of Fluctuating Temperature on Mortality, Stress, and Energy Reserves of Juvenile Coho Salmon. Transactions of the American Fisheries Society 115:52-59.
- Udey, L.R., J.L. Fryer, and K.S. Pilcher. 1975. Relation of water temperature to ceratomyxosis in rainbow trout (*Salmo gairdneri*) and coho salmon (*Oncorhynchus kisutch*). J. Fish. Res. Board Can. 32:1545-1551.
- U.S. Environmental Protection Agency (USEPA). 1986a. Ambient Water Quality Criteria for Dissolved Oxygen. Office of Water. EPA 440/5-86-003. 46pp.
- U.S. Environmental Protection Agency (USEPA). 1986b. Quality Criteria for Water. Office of Water. EPA 440/5-86-001. 477pp. Available at: http://epa.gov/waterscience/criteria/goldbook.pdf>. Website accessed August 12, 2005.
- U.S. Environmental Protection Agency (USEPA). 1999a. A review and synthesis of effects of alternation to the water temperature regime on freshwater life stages of salmonids, with special reference to Chinook salmon. Region 10, Seattle, WA.

- EPA 910-R-99-010. 279pp. Available at: http://www.critfc.org/tech/EPA/ report.htm>. Website accessed on June 11, 2004.
- U.S. Environmental Protection Agency (USEPA). 1999b. 1999 Update of Ambient Water Quality Criteria for Ammonia. Office of Water. EPA-822-R-99-014. 147pp.
- U.S. Environmental Protection Agency (USEPA). 2001. Issue Paper 5: Summary of technical literature examining the effects of temperature on salmonids. Region 10, Seattle, WA. EPA 910-D-01-005. 113pp. Available at: http://yosemite.epa.gov/R10/water.nsf>. Website accessed on July 2, 2004.
- U.S. Environmental Protection Agency (USEPA). 2003. EPA Region 10 Guidance for Pacific Northwest State and Tribal Water Quality Standards. Region 10, Seattle, WA. EPA 910-B-03-002. 49pp. Available at: http://www.epa.gov/r10earth/temperature.htm. Website accessed on June 23, 2004.
- Washington State Department of Ecology (WDOE). 2002. Evaluating Criteria for the Protection of Freshwater Aquatic Life in Washington's Surface Water Quality Standards: Dissolved Oxygen. Draft Discussion Paper and Literature Summary. Publication Number 00-10-071. 90pp.
- Wagner, E.J., T. Bosakowski, and S. Intelmann. 1997. Combined Effects of Temperature and High pH on Mortality and the Stress Response of Rainbow Trout after Stocking. Transactions of the American Fisheries Society. 126:985-998.
- Weiner G.S., C.B. Schreck, and H.W. Li 1986. Effects of Low pH on Reproduction of Rainbow Trout. Transactions of the American Fisheries Society. 115:75-82.
- Welsh, H.W., Jr., G.R. Hodgson, B.R. Harvey, and M.F. Roche. 2001. Distribution of juvenile coho salmon in relation to water temperatures in tributaries of the Mattole River, California. North American Journal of Fisheries Management 21:464-470.
- Whitmore, C.M., C.E. Warren, and P. Doudoroff. 1960. Avoidance reactions of salmonid and centrarchid fishes to low oxygen concentrations. Transactions of the American Fisheries Society. 89:17-26.