



MONITORING AND ANALYSIS PLAN

Second Statewide Survey of Bioaccumulation on the California Coast: 2024

Bioaccumulation Monitoring Program

Surface Water Ambient Monitoring Program

Version 1

December 2023



Table of Contents

| MONITORING AND ANALYSIS PLAN | 1 |
|--|----|
| Background | 3 |
| Monitoring Design Updates | 3 |
| References | 4 |
| Table 2. Long-term sport fish sampling schedule | 5 |
| Table 6. Target species by region for coastal waters and bays and harbors | 5 |
| Table 8. Target species, size ranges, and numbers to include in composites | 8 |
| Table 10. Parameters to be measured by the SWAMP labs | 12 |
| Figure A. Coastal zones to be sampled in 2024 | 13 |

Background

The Surface Water Ambient Monitoring Program (SWAMP) conducted an initial two-year statewide survey of bioaccumulation on the California coast in 2009-2010 (<u>Davis et al. 2012</u>). The work was conducted under the guidance of the Bioaccumulation Oversight Group (BOG), a SWAMP subcommittee. The BOG published a sampling and analysis plan for this screening study in 2009 (<u>BOG 2009</u>).

In 2018, SWAMP and the BOG began a second round of statewide sampling of the California coast. A sampling and analysis plan for this effort was published by the BOG in 2018 (BOG 2018). The original plan was to conduct this survey over a three-year period (2018-2020). The Southern California Bight was sampled as planned in 2018, in collaboration with the Southern California Bight Program. The results of this monitoring have been published (McLaughlin et al. 2020, 2021). Due to budgeting issues, contracting issues, and the pandemic of 2020, sampling of the rest of the coast has been spread over more years than originally anticipated. In addition to the Southern California Bight zones, some Central Coast zones were sampled in 2018. In 2019 Humboldt Bay and two zones near Santa Cruz were sampled. In 2020, 12 zones were sampled in the Central and North Coast regions, as documented in the cruise report for that year (MPSL 2020).

The remaining 18 zones in the Central and North Coast regions will be sampled in 2024. This monitoring and analysis plan documents the plans for completing the second round of coast sampling in 2024.

Monitoring Design Updates

The Safe to Eat Workgroup (STEW; formerly known as the BOG) has provided valuable input and feedback throughout the monitoring process, which has resulted in minor modifications to the monitoring design published in BOG (2018). This update to the original 2018 Monitoring Plan documents the modifications that have been made. All other aspects of the monitoring design match what was outlined in BOG (2018).

The tables and figures that have been updated from the original 2018 Monitoring Plan are provided below, following the numbering used in the original Monitoring Plan. Where changes were minor, revisions to the original tables are highlighted in yellow.

In addition to the updated tables, Figure A is included to provide a map of the 18 zones to be sampled in 2024.

As in the original design, the plan for individual contaminants is as follows.

- Mercury: All samples individuals in key indicator species, composites for others
- PCBs: Composites of 2 species per zone
- Selenium: Composites of all samples
- Archived samples can be analyzed for other analytes

References

- BOG. 2009. Sampling and Analysis Plan for a Screening Study of Bioaccumulation on the California Coast. Bioaccumulation Oversight Group, Surface Water Ambient Monitoring Program, State Water Resources Control Board, Sacramento, CA. https://www.waterboards.ca.gov/water_issues/programs/swamp/docs/bog_coast_fin_al_monplan_09.pdf
- BOG. 2018. Monitoring Plan for a Second Statewide Survey of Bioaccumulation on the California Coast. Bioaccumulation Oversight Group, Surface Water Ambient Monitoring Program, State Water Resources Control Board, Sacramento, CA. https://drive.google.com/file/d/1DlckuBLtgEZigNnfvRRxrv1UBDQ6zudV/view
- Davis, J.A., J.R.M. Ross, S.N. Bezalel, J.A. Hunt, A.R. Melwani, R.M. Allen, G. Ichikawa, A. Bonnema, W.A. Heim, D. Crane, S. Swenson, C. Lamerdin, M. Stephenson, and K. Schiff. 2012. Contaminants in Fish from the California Coast, 2009-2010: Summary Report on a Two-Year Screening Survey. A Report of the Surface Water Ambient Monitoring Program (SWAMP). California State Water Resources Control Board, Sacramento, CA. https://www.waterboards.ca.gov/water_issues/programs/swamp/docs/coast_study/bog2012may/coast2012report.pdf
- Klasing, S., W. Smith, L. Chumney. 2022. Protocol for Fish Sampling and Analysis to Support the Development of Fish Advisories in California. Office of Environmental Health Hazard Assessment, Sacramento, CA. https://oehha.ca.gov/media/downloads/fish/report/fishadvisorysamplinganalysisprotocolreport2022.pdf
- McLaughlin, K., K. Schiff, B. Du, J. Davis, A. Bonnema, G. Ichikawa, B. Jakl, and W. Heim. 2020. Southern California Bight 2018 Regional Monitoring Program: Volume V. Contaminant Bioaccumulation in Edible Sport Fish Tissue. Technical Report 1155. Southern California Coastal Water Research Project. Costa Mesa, CA. https://ftp.sccwrp.org/pub/download/DOCUMENTS/TechnicalReports/1155 B18Bioa ccumulation.pdf
- McLaughlin, K., J.A. Davis, A. Bonnema, B. Du, G. Ichikawa, W. Jakl, W. Heim, and K. Schiff. 2021. Regional assessment of contaminant bioaccumulation in sport fish tissue in the Southern California Bight, USA. Marine Pollution Bulletin, 172, p.112798.
- MPSL. 2020. Cruise Report for the Surface Water Ambient Monitoring Program (SWAMP) A Second Statewide Survey of Bioaccumulation on the California Coast (Year 2 of 3 Years). Marine Pollution Studies Laboratory, Moss Landing Marine Laboratories, Moss Landing, CA.
 - https://drive.google.com/file/d/1hAxmw1kJPWK6OELqbTmUrSmU79XskdZ4/view

Table 2. Long-term sport fish sampling schedule

At the time of publication of this 2024 sampling plan update, the long-term sampling schedule for the SWAMP Bioaccumulation Monitoring Program is under discussion and to be determined.

Table 6. Target species by region for coastal waters and bays and harbors

Species in italics are those that will be analyzed as individuals for mercury as well as composited for other analytes. If the target species to be analyzed as individuals for mercury are not available, substitutions will be made.

Asterisks (*) indicate species that were in the top five in catch based on RecFIN data for each habitat by region combination.

Bold indicates primary target species for the Bight Program. Based on results of the San Diego Bay Consumption Study, primary targets in San Diego Bay are chub mackerel, spotted sand bass, halibut, and topsmelt.

Entries highlighted in yellow are additions to the list based on input from the Department of Public Health, discussed at the STEW meeting on October 18, 2023 (the changes to the list for Southern California are noted for future monitoring in this region).

Coast (>3 miles)

| Targe Species Type (Primary or Secondary) | South | Central | North |
|---|------------------|-------------------|-----------------|
| Primary | Kelp Bass | | |
| Primary | Barred Sand Bass | | |
| Primary | Gopher Rockfish | Gopher Rockfish | Gopher Rockfish |
| Primary | | Brown Rockfish | Brown Rockfish |
| Primary | | Black Rockfish* | Black Rockfish* |
| Primary | | Blue Rockfish* | Blue Rockfish* |
| Primary | Lingcod | Lingcod* | Lingcod* |
| Primary | | Cabezon | Cabezon |
| Primary | | Salmon | Salmon |
| Primary | Chub Mackerel* | | |
| Primary | White Croaker | White Croaker | |
| Primary | | Rainbow Surfperch | |
| Secondary | Scorpionfish | | |

| Targe Species Type (Primary or Secondary) | South | Central | North |
|---|------------------------------------|----------------------|----------------------|
| Secondary | Blue Rockfish | | |
| Secondary | Brown Rockfish | | |
| Secondary | Copper Rockfish | Copper Rockfish | Copper Rockfish |
| Secondary | Vermilion Rockfish | Vermilion Rockfish | Vermilion Rockfish |
| Secondary | | Olive Rockfish | Olive Rockfish |
| Secondary | | Yellowtail Rockfish* | Yellowtail Rockfish* |
| Secondary | Barred Surfperch* | Barred Surfperch* | Barred Surfperch* |
| Secondary | Walleye Surfperch | Walleye Surfperch | Walleye Surfperch |
| Secondary | | Redtail Surfperch | Redtail Surfperch |
| Secondary | California Halibut | California Halibut | California Halibut |
| Secondary | Yellowfin Croaker | | |
| Secondary | California Sheephead (OEHHA) | | |
| Secondary | Halfmoon/Opaleye (OEHHA) | | |
| Secondary | Kelp Greenling (OEHHA) | | |
| Secondary | Pacific Halibut (OEHHA) | | |
| Secondary | | Jacksmelt* | |
| Secondary | California Corbina (DPH) | | |
| Secondary | Bocaccio (DPH) | | |
| Secondary | | White Seabass | White Seabass |
| Secondary | | Chub Mackerel | Chub Mackerel |

Bays and Harbors

| Targe Species Type (Primary or Secondary) | South | Central | North | | |
|---|-------------------|----------------|-------|--|--|
| Primary | Kelp Bass | | | | |
| Primary | | Brown Rockfish | | | |
| Primary | Spotted Sand Bass | | | | |

| Targe Species Type (Primary or Secondary) | South | Central | North |
|---|-----------------------------|---------------------|---------------------|
| Primary | Shiner Surfperch | Shiner Surfperch | Shiner Surfperch |
| Primary | | | White Surfperch |
| Primary | | Leopard Shark | Leopard Shark |
| Primary | | California Halibut* | California Halibut* |
| Primary | Jacksmelt* | Jacksmelt* | Jacksmelt* |
| Primary | White Croaker | White Croaker | White Croaker |
| Primary | Chub Mackerel* | | |
| Primary | | Striped Bass* | |
| Secondary | Barred Sand Bass | | |
| Secondary | Scorpionfish | | |
| Secondary | | Chub Mackerel* | Chub Mackerel* |
| Secondary | | Black Rockfish | Black Rockfish |
| Secondary | Spotfin Croaker* | | |
| Secondary | Yellowfin Croaker* | | |
| Secondary | White Surfperch | | |
| Secondary | | Black Perch | Black Perch |
| Secondary | California Halibut | | |
| Secondary | Leopard Shark | | |
| Secondary | Gray Smoothound | | |
| Secondary | Brown Smoothound | Brown Smoothound | Brown Smoothound |
| Secondary | Spiny Dogfish | Spiny Dogfish | Spiny Dogfish |
| Secondary | Topsmelt | | |
| Secondary | | Bat Ray | Bat Ray |
| Secondary | Halfmoon/Opaleye (OEHHA) | | |
| Secondary | | | Redtail Surfperch |
| Secondary | | | Walleye Surperch |

Table 8. Target species, size ranges, and numbers to include in composites

Items revised from the original 2018 Monitoring Plan are highlighted in yellow. Size range changes are based on guidance published in 2022 by OEHHA (<u>Klasing et al. 2022</u>).

| Group | Species | Primary or Secondary Target (P or S) | Mercury in Individuals or Composites (I or C) | # fish in C | Min Legal Size (in) | Min Legal Size (mm) | Median Size 2009-2010* (mm) | Targeted Size Range Individuals (mm) | Size Range Composites (mm) |
|----------|---------------------|---|---|-------------------|------------------------------|------------------------------|--------------------------------------|---|----------------------------------|
| Rockfish | Kelp Bass | Р | I,C | 5 | 14 | 356 | 316 | 3X(156-255) 3X(256-355) 5X(>356) | >356 |
| Rockfish | Barred Sandbass | P,S | I,C | 5 | 14 | 356 | 346 | 3X(156-255) 3X(256-355) 5X(>356) | >356 |
| Rockfish | Spotted Sandbass | Р | I,C | 5 | 14 | 356 | 327 | 3X(156-255) 3X(256-355) 5X(>356) | >356 |
| Rockfish | Gopher Rockfish | Р | I | | | | 281 | 3X(141-191) 3X(191-241) 5X(>241) | >135 |
| Rockfish | Blue Rockfish | P,S | С | 5 | | | 293 | | >200 |

| Group | Species | Primary or Secondary Target (P or S) | Mercury in Individuals or Composites (I or C) | # fish in C | Min Legal Size (in) | Min Legal Size (mm) | Median Size 2009-2010* (mm) | Targeted Size Range Individuals (mm) | Size Range Composites (mm) |
|----------|------------------------|---|---|-------------------|------------------------------|------------------------------|--------------------------------------|---|----------------------------------|
| Rockfish | Black Rockfish | P,S | I | | | | 380 | 3X(156-255) 3X(256-354) 5X(>355) | >355 |
| Rockfish | Scorpionfish | S | С | 5 | 10 | 254 | 290 | | >254 |
| Rockfish | Olive Rockfish | S | С | 5 | | | 322 | | >320 |
| Rockfish | Brown Rockfish | P,S | I | | | | 302 | 3X(150-199) 2X(200-249) 5X(>250) | >250 |
| Rockfish | Copper Rockfish | S | С | 5 | | | 411 | | >320 |
| Rockfish | Vermilion Rockfish | S | C | <mark>5</mark> | | | | | >350 |
| Rockfish | Bocaccio | S | C | <mark>5</mark> | | | | | <mark>??</mark> |
| Rockfish | Yellowtail Rockfish | S | C | <mark>5</mark> | | | <mark>313</mark> | | >290* |
| Lingcod | | P,S | I | | 22 | 559 | 682 | 3X(359-458) 2X(459-558) 5X(>559) | >559 |
| Croaker | White Croaker | P,S | С | 5 | | | 220 | | >150 |
| Croaker | Yellowfin Croaker | S | С | 5 | | | 195 | | >167 |

| Group | Species | Primary or Secondary Target (P or S) | Mercury in Individuals or Composites (I or C) | # fish in C | Min Legal Size (in) | Min Legal Size (mm) | Median Size 2009-2010* (mm) | Targeted Size Range Individuals (mm) | Size Range Composites (mm) |
|-------------------|-----------------------|---|---|-------------------|------------------------------|------------------------------|--------------------------------------|---|----------------------------------|
| Croaker | Spotfin Croaker | S | С | 5 | | | 221 | | >189 |
| Croaker | California Corbina | S | C | <mark>5</mark> | | | | | >250 |
| Croaker | White Seabass | S | C | <mark>5</mark> | | | | | <mark>??</mark> |
| Chinook Salmon | | Р | С | | 24 | Check regs | | Not included in 2024 | |
| Surfperch | Barred | S | С | 5 | | | 186 | | <mark>>160</mark> |
| Surfperch | Redtail | S | С | 5 | 10.5 | 267 | No data | | >267 |
| Surfperch | Shiner | Р | С | 20 | | | 110 | | >100 |
| Surfperch | Walleye | S | С | 5 | | | No data | | <mark>>115</mark> |
| Surfperch | Black | S | С | 5 | | | 232 | | <mark>>160</mark> |
| Surfperch | Rainbow | Р | С | 5 | | | 280 | | <mark>>125</mark> |
| Surfperch | White | P,S | С | 5 | | | 202 | | <mark>>125</mark> |
| Smelt | Jacksmelt | P,S | С | 5 | | | 265 | | >227 |
| Smelt | Topsmelt | S | С | 5 | | | 128 | | <mark>>150</mark> |
| Chub Mackerel | | Р | С | 5 | | | 240 | | >260 |

| Group | Species | Primary or Secondary Target (P or S) | Mercury in Individuals or Composites (I or C) | # fish in C | Min Legal Size (in) | Min Legal Size (mm) | Median Size 2009-2010* (mm) | Targeted Size Range Individuals (mm) | Size Range Composites (mm) |
|-------------------|-----------------------|---|---|-------------------|------------------------------|------------------------------|--------------------------------------|--|----------------------------------|
| Shark | Leopard Shark | Р | l | | 36 | 914 | 1238 | 3X(914- 1074) 4X(1074- 1234) 3X(>1234) | >914 |
| Shark | Spiny Dogfish | S | С | 3 | | | 1011 | | >867 |
| Shark | Brown Smoothhound | S | С | 3 | | | 978 | | <mark>>610</mark> |
| Shark | Gray Smoothound | S | С | 3 | | | 630 | | <mark>>580</mark> |
| Ray | Bat Ray | S | С | 3 | | | 405 | | >347 |
| Halibut | California Halibut | P,S | С | 3 | | <mark>559</mark> | 670 | | >559 |
| Halibut | Pacific Halibut | S | С | 3 | | | No data | | ?? |
| Cabezon | | P,S | I | | 15 | 381 | 467 | 3X(281-330) 3X(331-380) 5X(>381) | >381 |
| Halfmoon | | S | С | 5 | | | No data | | |
| Opaleye | | S | С | 5 | | | 221 | | >189 |
| Kelp Greenling | | S | С | 5 | 12 | 305 | 360 | | >309 |

| Group | Species | Primary or Secondary Target (P or S) | Mercury in Individuals or Composites (I or C) | # fish in C | Min Legal Size (in) | Min Legal Size (mm) | Median Size 2009-2010* (mm) | Targeted Size Range Individuals (mm) | Size Range Composites (mm) |
|-------------------------|---------|---|---|-------------------|------------------------------|------------------------------|--------------------------------------|---|----------------------------------|
| California Sheephead | | S | С | 5 | 12 | 305 | No data | | >305 |
| Striped Bass | | S | Ī | 3 | 18 | <mark>457</mark> | | 3X(257-356) 3X(357-456) 5X(>457) | >457 |

^{*} based on minimum size caught in the 2009-2010 statewide survey.

Table 10. Parameters to be measured by the SWAMP labs

In the 2018 Plan, 53 PCB congeners were included on the analyte list in Table 10. The number of PCB congeners analyzed has been increasing over the years. In 2024 the full suite of 209 congeners will be analyzed. The sums of PCBs reported will include all 209 congeners, with "not detected" values for individual congeners set to zero. For rigorous comparisons with past data a subset of common congeners can be used.

Figure A. Coastal zones to be sampled in 2024

